

# Coarse approximation in convex analysis

Grigory Ivanov



Conference on Convex Geometry and Geometric Probability,  
Paris Lodron University Salzburg

September 26, 2023

# Disclaimer

Caveat emptor (Let the buyer beware)

# Outline

From “Sparse” to “Coarse”

Helly-type results

Open problems

# “Sparse” approximation

Consider

$$\text{Id}_d = \sum_{i=1}^m c_i u_i \otimes u_i = \sum_{i=1}^m c_i u_i u_i^T,$$

where  $c_1, \dots, c_m$  are positive weights;  $u_1, \dots, u_m$  are unit vectors in  $\mathbb{R}^d$ .

# “Sparse” approximation

Consider

$$\text{Id}_d = \sum_{i=1}^m c_i u_i \otimes u_i = \sum_{i=1}^m c_i u_i u_i^T,$$

where  $c_1, \dots, c_m$  are positive weights;  $u_1, \dots, u_m$  are unit vectors in  $\mathbb{R}^d$ .

**Problem:**

For a given  $\varepsilon$ , find a **smallest** index set  $J$  such that

$$\frac{\text{Id}_d}{1 + \varepsilon} \prec \sum_{j \in J} \tilde{c}_j u_j \otimes u_j \prec (1 + \varepsilon) \text{Id}_d$$

for some positive weights  $\{\tilde{c}_j\}_{j \in J}$

Intuition says  $d^2 \dots$

Intuition says  $d^2 \dots$

Bourgain'95

$$k \leq C(\varepsilon) d (\ln d)^3$$

Intuition says  $d^2$ ...

Bourgain'95

$$k \leq C(\varepsilon)d(\ln d)^3$$

Rudelson'99

$$k \leq C \frac{d \ln d}{\varepsilon^2}$$



Intuition says  $d^2$ ...

Bourgain'95

$$k \leq C(\varepsilon)d(\ln d)^3$$

Rudelson'99

$$k \leq C \frac{d \ln d}{\varepsilon^2}$$

Breakthrough! BSS'14, MSS'15, FY'17

$$k \leq C \frac{d}{\varepsilon^2}$$

Intuition says  $d^2$ ...

Bourgain'95

$$k \leq C(\varepsilon)d(\ln d)^3$$

Rudelson'99

$$k \leq C \frac{d \ln d}{\varepsilon^2}$$

Breakthrough! BSS'14, MSS'15, FY'17

$$k \leq C \frac{d}{\varepsilon^2}$$

# What are the first steps?

# What are the first steps?

Consider  $\text{Id}_d = \sum_{i=1}^m c_i u_i \otimes u_i$

# What are the first steps?

Consider  $\text{Id}_d = \sum_{i=1}^m c_i u_i \otimes u_i$

## Qualitative Question:

What is the size of the smallest index set  $J$  such that

$$\frac{\text{Id}_d}{1 + \lambda_d} \prec \sum_{j \in J} \tilde{c}_j u_j \otimes u_j \prec (1 + \lambda_d) \text{Id}_d$$

for some positive weights  $\{\tilde{c}_j\}_{j \in J}$  and **some**  $\lambda_d$ .

# What are the first steps?

Consider  $\text{Id}_d = \sum_{i=1}^m c_i u_i \otimes u_i$

## Qualitative Question:

What is the size of the smallest index set  $J$  such that

$$\frac{\text{Id}_d}{1 + \lambda_d} \prec \sum_{j \in J} \tilde{c}_j u_j \otimes u_j \prec (1 + \lambda_d) \text{Id}_d$$

for some positive weights  $\{\tilde{c}_j\}_{j \in J}$  and **some**  $\lambda_d$ .

**Answer:** Trivial, the size is  $d$ .

# What the first steps give us quantitatively?

## Quantitative Question:

What is the smallest  $\lambda_d$  such that

# What the first steps give us quantitatively?

## Quantitative Question:

What is the smallest  $\lambda_d$  such that for **any** “John’s decompositions”

$$\text{Id}_d = \sum_{i=1}^m c_i u_i \otimes u_i$$



# What the first steps give us quantitatively?

## Quantitative Question:

What is the smallest  $\lambda_d$  such that for **any** “John’s decompositions”  
 $\text{Id}_d = \sum_{i=1}^m c_i u_i \otimes u_i$  one can find an index set  $J$  of size  $d$  and

# What the first steps give us quantitatively?

## Quantitative Question:

What is the smallest  $\lambda_d$  such that for **any** “John’s decompositions”  $\text{Id}_d = \sum_{i=1}^m c_i u_i \otimes u_i$  one can find an index set  $J$  of size  $d$  and positive weights  $\{\tilde{c}_j\}_{j \in J}$  satisfying

$$\frac{\text{Id}_d}{1 + \lambda_d} \prec \sum_{j \in J} \tilde{c}_j u_j \otimes u_j \prec (1 + \lambda_d) \text{Id}_d$$

# What the first steps give us quantitatively?

## Quantitative Question:

What is the smallest  $\lambda_d$  such that for **any** “John’s decompositions”  $\text{Id}_d = \sum_{i=1}^m c_i u_i \otimes u_i$  one can find an index set  $J$  of size  $d$  and positive weights  $\{\tilde{c}_j\}_{j \in J}$  satisfying

$$\frac{\text{Id}_d}{1 + \lambda_d} \prec \sum_{j \in J} \tilde{c}_j u_j \otimes u_j \prec (1 + \lambda_d) \text{Id}_d$$

Answer

# What the first steps give us quantitatively?

## Quantitative Question:

What is the smallest  $\lambda_d$  such that for **any** “John’s decompositions”  $\text{Id}_d = \sum_{i=1}^m c_i u_i \otimes u_i$  one can find an index set  $J$  of size  $d$  and positive weights  $\{\tilde{c}_j\}_{j \in J}$  satisfying

$$\frac{\text{Id}_d}{1 + \lambda_d} \prec \sum_{j \in J} \tilde{c}_j u_j \otimes u_j \prec (1 + \lambda_d) \text{Id}_d$$

Answer or not:

# What the first steps give us quantitatively?

## Quantitative Question:

What is the smallest  $\lambda_d$  such that for **any** “John’s decompositions”  $\text{Id}_d = \sum_{i=1}^m c_i u_i \otimes u_i$  one can find an index set  $J$  of size  $d$  and positive weights  $\{\tilde{c}_j\}_{j \in J}$  satisfying


$$\frac{\text{Id}_d}{1 + \lambda_d} \prec \sum_{j \in J} \tilde{c}_j u_j \otimes u_j \prec (1 + \lambda_d) \text{Id}_d$$

Answer or not:

... take the vectors by the “Dvoretzky–Rogers lemma”...


# John ellipsoid

1.  $K$  is a convex body in  $\mathbb{R}^d$



# John ellipsoid


1.  $K$  is a convex body in  $\mathbb{R}^d$
2. The *John ellipsoid*  $J_K$  of  $K$  is the maximal volume ellipsoid contained within  $K$



# John ellipsoid

1.  $K$  is a convex body in  $\mathbb{R}^d$
2. The *John ellipsoid*  $J_K$  of  $K$  is the maximal volume ellipsoid contained within  $K$
3. Let  $0$  be the center of  $J_K$ . Then

$$J_K \subset K \subset d \cdot J_K$$





# John condition

$K \subset \mathbb{R}^d$  convex body.

## John '48 (+ Ball'92)

Assume  $\mathbf{B}^d \subseteq K$ . TFAE:

1.  $\mathbf{B}^d$  is the maximum volume ellipsoid contained within  $K$
2. there are contact points  $u_1, \dots, u_m \in \text{bd } \mathbf{B}^d \cap \text{bd } K$  and positive weights  $c_1, \dots, c_m$  such that

$$\sum_{i=1}^m c_i u_i = 0 \quad \text{and} \quad \sum_{i=1}^m c_i u_i \otimes u_i = \text{Id}_d$$

# Outline

From “Sparse” to “Coarse”

Helly-type results

Open problems

# Classical result

Helly'1923\*

If the intersection of every  $d + 1$  sets of a finite family of convex subsets of  $\mathbb{R}^d$  is nonempty, then the whole family has a nonempty intersection.

# Classical result

Helly'1923\*

If the intersection of every  $d + 1$  sets of a finite family of convex subsets of  $\mathbb{R}^d$  is nonempty, then the whole family has a nonempty intersection.

Reduction assumption: All sets in our family are half-spaces!

# Classical result

Helly'1923\*

If the intersection of every  $d + 1$  sets of a finite family of convex subsets of  $\mathbb{R}^d$  is nonempty, then the whole family has a nonempty intersection.

Reduction assumption: All sets in our family are half-spaces!

## Notation

For  $v \in \mathbb{R}^d \setminus \{0\}$ ,


$$H_v = \{p \in \mathbb{R}^d : \langle p, v \rangle \leq 1\}$$

# Coarse approximation

Reduction assumption: All sets in our family are half-spaces!


# Coarse approximation

Reduction assumption: All sets in our family are half-spaces!



# Coarse approximation


Reduction assumption: All sets in our family are half-spaces!






# Coarse approximation

Reduction assumption: All sets in our family are half-spaces!



# Coarse approximation

Reduction assumption: All sets in our family are half-spaces!



# Coarse approximation

Reduction assumption: All sets in our family are half-spaces!

## Qualitative Question:

How many half-spaces (facets) must we retain to ensure that the intersection forms a bounded polytope?

# Coarse approximation

Reduction assumption: All sets in our family are half-spaces!

## Quantitative Question:

How many half-spaces (facets) must we retain to ensure that the intersection forms a bounded polytope?

Helly-number: Sometimes we need  $2d$  half-spaces — Cube!

# Coarse approximation

Reduction assumption: All sets in our family are half-spaces!

## Quantitative Question:

How many half-spaces (facets) must we retain to ensure that the intersection forms a bounded polytope?

Helly-number: Sometimes we need  $2d$  half-spaces — Cube!

## Quantitative Question:

Given a convex polytope  $P$  in  $\mathbb{R}^d$ , can one choose at most  $m \leq 2d$  facets of  $P$  in such a way that the volume of the intersection of corresponding half-spaces is at most  $\nu_d \text{vol}_d P$ .

# Results

Bárány, Katchalski, Pach'82

Let  $\mathcal{F}$  be a finite family of convex subsets of  $\mathbb{R}^d$ .

Then one can find at most  $m \leq 2d$  sets  $F_1, \dots, F_m$  of  $\mathcal{F}$  satisfying

$$\text{vol}_d(F_1 \cap \dots \cap F_m) \leq \nu_d \text{vol}_d \bigcap \mathcal{F}$$

# Results

Bárány, Katchalski, Pach'82

Let  $\mathcal{F}$  be a finite family of convex subsets of  $\mathbb{R}^d$ .

Then one can find at most  $m \leq 2d$  sets  $F_1, \dots, F_m$  of  $\mathcal{F}$  satisfying

$$\text{vol}_d(F_1 \cap \dots \cap F_m) \leq \nu_d \text{vol}_d \bigcap \mathcal{F} \quad (\nu_d \approx d^{2d^2}).$$

# Results

## Bárány, Katchalski, Pach'82

Let  $\mathcal{F}$  be a finite family of convex subsets of  $\mathbb{R}^d$ .

Then one can find at most  $m \leq 2d$  sets  $F_1, \dots, F_m$  of  $\mathcal{F}$  satisfying

$$\text{vol}_d(F_1 \cap \dots \cap F_m) \leq \nu_d \text{vol}_d \bigcap \mathcal{F} \quad (\nu_d \approx d^{2d^2}).$$

## Bárány, Katchalski, Pach'82

Let  $\mathcal{F}$  be a finite family of convex subsets of  $\mathbb{R}^d$ .

Then one can find at most  $m \leq 2d$  sets  $F_1, \dots, F_m$  of  $\mathcal{F}$  satisfying

$$\text{diam}(F_1 \cap \dots \cap F_m) \leq \eta_d \text{diam} \bigcap \mathcal{F}$$



## Bárány, Katchalski, Pach'82

Let  $\mathcal{F}$  be a finite family of convex subsets of  $\mathbb{R}^d$ .

Then one can find at most  $m \leq 2d$  sets  $F_1, \dots, F_m$  of  $\mathcal{F}$  satisfying

$$\text{vol}_d(F_1 \cap \dots \cap F_m) \leq \nu_d \text{vol}_d \bigcap \mathcal{F} \quad (\nu_d \approx d^{2d^2}).$$

## Bárány, Katchalski, Pach'82

Let  $\mathcal{F}$  be a finite family of convex subsets of  $\mathbb{R}^d$ .

Then one can find at most  $m \leq 2d$  sets  $F_1, \dots, F_m$  of  $\mathcal{F}$  satisfying

$$\text{diam}(F_1 \cap \dots \cap F_m) \leq \eta_d \text{diam} \bigcap \mathcal{F} \quad (\eta_d \approx d^{d/2})$$

## Naszódi'16

Let  $\mathcal{F}$  be a finite family of convex subsets of  $\mathbb{R}^d$ .

Then one can find at most  $m \leq 2d$  sets  $F_1, \dots, F_m$  of  $\mathcal{F}$  satisfying

$$\text{vol}_d(F_1 \cap \dots \cap F_m) \leq \nu_d \text{vol}_d \bigcap \mathcal{F} \quad \left( \nu_d \approx d^{10d} \right).$$

$$\nu_d > d^{d/2}$$

# Results

## Naszódi'16

Let  $\mathcal{F}$  be a finite family of convex subsets of  $\mathbb{R}^d$ .

Then one can find at most  $m \leq 2d$  sets  $F_1, \dots, F_m$  of  $\mathcal{F}$  satisfying

$$\text{vol}_d(F_1 \cap \dots \cap F_m) \leq \nu_d \text{vol}_d \bigcap \mathcal{F} \quad (\nu_d \approx d^{10d}).$$

$$\nu_d > d^{d/2}$$

## Brazitikos'18

Let  $\mathcal{F}$  be a finite family of convex subsets of  $\mathbb{R}^d$ .

Then one can find at most  $m \leq 2d$  sets  $F_1, \dots, F_m$  of  $\mathcal{F}$  satisfying

$$\text{diam}(F_1 \cap \dots \cap F_m) \leq \eta_d \text{diam} \bigcap \mathcal{F} \quad (\eta_d \approx d^{11/2}).$$

Extensive use of the John ellipsoid and the Dvoretzky–Rogers...!

# Containment in a homothet

Do we need the John ellipsoid in the proofs?

# Containment in a homothet

Do we need the John ellipsoid in the proofs?

**NO!**

# Containment in a homothet

What do we really use?

# Containment in a homothet

What do we really use?

1. Duality:

## Polar set

The polar  $S^\circ$  of a set  $S \subset \mathbb{R}^d$  is the set

$$S^\circ = \left\{ p \in \mathbb{R}^d : \quad \langle p, x \rangle \leq 1 \quad \text{for all } x \in S \right\}$$

# Containment in a homothet

What do we really use?

1. Duality:

$$K = \bigcap_{i \in I} H_{v_i} \quad \Longleftrightarrow \quad K^\circ = \text{conv}\{v_i : i \in I\}$$



# Containment in a homothet

What do we really use?

1. Duality:

$$K = \bigcap_{i \in I} H_{v_i} \quad \Longleftrightarrow \quad K^\circ = \text{conv}\{v_i : i \in I\}$$

$$K \subset \bigcap_{i \in J \cup I} H_{v_i} \quad \Longleftrightarrow \quad K^\circ \supset \text{conv}\{v_i : i \in J\}$$

# Containment in a homothet

What do we really use?

1. Duality:

$$K = \bigcap_{i \in I} H_{v_i} \quad \Longleftrightarrow \quad K^\circ = \text{conv}\{v_i : i \in I\}$$

$$K \subset \bigcap_{i \in J \cup I} H_{v_i} \quad \Longleftrightarrow \quad K^\circ \supset \text{conv}\{v_i : i \in J\}$$

2. Double-sided inclusion:

$$\bigwedge \text{conv}\{v_i : i \in J\} \supset K^\circ \supset \text{conv}\{v_i : i \in J\}$$

# Containment in a homothet

What do we really use?

1. Duality:

$$K = \bigcap_{i \in I} H_{v_i} \quad \Longleftrightarrow \quad K^\circ = \text{conv}\{v_i : i \in I\}$$

$$K \subset \bigcap_{i \in J \cup I} H_{v_i} \quad \Longleftrightarrow \quad K^\circ \supset \text{conv}\{v_i : i \in J\}$$

2. Double-sided inclusion:

$$|\Lambda| \text{conv}\{v_i : i \in J\} \supset K^\circ \supset \text{conv}\{v_i : i \in J\}$$

$$\Downarrow$$

$$\text{vol}_d \bigcap_{i \in J \cup I} H_{v_i} \leq |\Lambda|^d \text{vol}_d K \quad \text{and} \quad \text{diam} \bigcap_{i \in J \cup I} H_{v_i} \leq |\Lambda| \text{diam} K$$

# Containment in a homothet

What do we really use?

1. Duality:

$$K = \bigcap_{i \in I} H_{v_i} \quad \Longleftrightarrow \quad K^\circ = \text{conv}\{v_i : i \in I\}$$

$$K \subset \bigcap_{i \in J \cup I} H_{v_i} \quad \Longleftrightarrow \quad K^\circ \supset \text{conv}\{v_i : i \in J\}$$

2. Double-sided inclusion:

$$\bigwedge \text{conv}\{v_i : i \in J\} \supset K^\circ \supset \text{conv}\{v_i : i \in J\}$$

3. “Good center”  $c$ :

$$K - c \subset -d(K - c)$$

# Containment in a homothet

What do we really use?

1. Duality:

$$K - \mathbf{c} = \bigcap_{i \in I} H_{v_i} \quad \Longleftrightarrow \quad (K - \mathbf{c})^\circ = \text{conv}\{v_i : i \in I\}$$

$$K - \mathbf{c} \subset \bigcap_{i \in J \subset I} H_{v_i} \quad \Longleftrightarrow \quad (K - \mathbf{c})^\circ \supset \text{conv}\{v_i : i \in J\}$$

2. Double-sided inclusion:

$$\bigwedge \text{conv}\{v_i : i \in J\} \supset (K - \mathbf{c})^\circ \supset \text{conv}\{v_i : i \in J\}$$

3. “Good center”  $\mathbf{c}$ :

$$K - \mathbf{c} \subset -\mathbf{d}(K - \mathbf{c})$$

The center of the John ellipsoid is a good center, but there are many others!

# Containment in a homothet: Double-sided inclusion

Naszódi and I.'22

Let  $S \subset \mathbb{R}^d$  be such that  $\text{conv } S \subset -d \text{ conv } S$ . Then there are  $m \leq 2d$  and  $v_1, \dots, v_m \in S$  such that

$$\text{conv } S \subset \Lambda \text{ conv}\{v_1, \dots, v_m\},$$

where  $\Lambda = -15d^3$ .

# Containment in a homothet: Double-sided inclusion

A-HAK lemma [Almendra–Hernández, Ambrus, Kendall'22]

Let  $S \subset \mathbb{R}^d$  be such that  $\text{conv } S \subset -d \text{ conv } S$ . Then there are  $m \leq 2d$  and  $v_1, \dots, v_m \in S$  such that

$$\text{conv } S \subset \Lambda \text{ conv}\{v_1, \dots, v_m\},$$

where  $\Lambda = -3d^2$ .

# Containment in a homothet: Double-sided inclusion

A-HAK lemma [Almendra–Hernández, Ambrus, Kendall'22]

Let  $S \subset \mathbb{R}^d$  be such that  $\text{conv } S \subset -d \text{ conv } S$ . Then there are  $m \leq 2d$  and  $v_1, \dots, v_m \in S$  such that

$$\text{conv } S \subset \Lambda \text{ conv}\{v_1, \dots, v_m\},$$


where  $\Lambda = -3d^2$ .

$v_1, \dots, v_d$  form a maximum volume  $d$ -simplex, i.e.


$$\max_{w_1, \dots, w_d \in S} \text{vol}_d \text{conv}\{0, w_1, \dots, w_d\} = \text{vol}_d \text{conv}\{0, v_1, \dots, v_d\}$$



# Containment in a homothet: Double-sided inclusion




# Containment in a homothet: Double-sided inclusion




# Containment in a homothet: Double-sided inclusion

$$(K - c)^0 \subset H_{v_2} \cap H_{-v_2}$$




# Containment in a homothet: Double-sided inclusion




# Containment in a homothet: Double-sided inclusion


$$(K - \mathbf{c})^0 \subset \bigcap_{i=1}^d (H_{v_i} \cap H_{-v_i})$$



# Containment in a homothet: Double-sided inclusion




# Containment in a homothet: Double-sided inclusion




## Containment in a homothet: Double-sided inclusion

$$n = \frac{V_1 + \dots + V_d}{d}$$





# Containment in a homothet: Double-sided inclusion



# Outline

From “Sparse” to “Coarse”

Helly-type results

Open problems

# Extremal arrangement of half-spaces

## Conjecture

Let  $\{u_1, \dots, u_{2d}\}$  be unit vectors in  $\mathbb{R}^d$ . Then there is a point in the set

$$\bigcap_{i=1}^{2d} \{x \in \mathbb{R}^d : \langle u_i, x \rangle \leq 1\}$$

with norm  $\sqrt{d}$ .

# Extremal arrangement of half-spaces

## Conjecture

Let  $\{u_1, \dots, u_{2d}\}$  be unit vectors in  $\mathbb{R}^d$ . Then there is a point in the set

$$\bigcap_{i=1}^{2d} \{x \in \mathbb{R}^d : \langle u_i, x \rangle \leq 1\}$$

with norm  $\sqrt{d}$ .

## Follows from a theorem of Ball and Prodromou'09

Let  $u_1, \dots, u_{2d}$  be unit vectors in  $\mathbb{R}^d$ . Then there is a point in the set

$$\bigcap_{i=1}^{2d} \{x \in \mathbb{R}^d : |\langle u_i, x \rangle| \leq 1\}$$

with norm  $\sqrt{\frac{d}{2}}$ .

## Good center: Macbeath point

### Macbeath region

Take a point  $p \in K \subset \mathbb{R}^d$ , the Macbeath region  $\mathcal{M}(K, p)$  is the set

$$\mathcal{M}(K, p) = K \cap (-K + 2p)$$

## Good center: Macbeath point

### Macbeath region

Take a point  $p \in K \subset \mathbb{R}^d$ , the Macbeath region  $\mathcal{M}(K, p)$  is the set

$$\mathcal{M}(K, p) = K \cap (-K + 2p)$$

### Macbeath region

Take a point  $p$  and a set  $K$ , the Macbeath region  $\mathcal{M}(K, p)$  is the set

$$\mathcal{M}(K, p) = K \cap (-K + 2p)$$

## Good center: Macbeath point

### Macbeath region

Take a point  $p \in K \subset \mathbb{R}^d$ , the Macbeath region  $\mathcal{M}(K, p)$  is the set

$$\mathcal{M}(K, p) = K \cap (-K + 2p)$$

### Proposition

Let  $K$  be a convex body in  $\mathbb{R}^d$ . There is a unique point  $p$  maximizing the volume of the Macbeath region  $\mathcal{M}(K, p)$ .

## Good center: Macbeath point

### Macbeath region

Take a point  $p \in K \subset \mathbb{R}^d$ , the Macbeath region  $\mathcal{M}(K, p)$  is the set

$$\mathcal{M}(K, p) = K \cap (-K + 2p)$$

### Proposition

Let  $K$  be a convex body in  $\mathbb{R}^d$ . There is a unique point  $p$  maximizing the volume of the Macbeath region  $\mathcal{M}(K, p)$ .

### Macbeath point

The Macbeath point of  $K$  is the unique maximizer of  $\text{vol}_d \mathcal{M}(K, p)$ .



## Good center: Macbeath point

### Macbeath region

Take a point  $p \in K \subset \mathbb{R}^d$ , the Macbeath region  $\mathcal{M}(K, p)$  is the set

$$\mathcal{M}(K, p) = K \cap (-K + 2p)$$

### Macbeath point

The Macbeath point of  $K$  is the unique maximizer of  $\text{vol}_d \mathcal{M}(K, p)$ .

### Question:

Let  $p$  be the Macbeath point of a convex body  $K \subset \mathbb{R}^d$ . Is it true that

$$K - p \subset -d(K - p)?$$

# Colorful versions

Qualitative result. Puzzle:

Let finite families  $\mathcal{F}_1, \dots, \mathcal{F}_{2d}$  of convex subsets of  $\mathbb{R}^d$  be such that  $\text{vol}_d \cap \mathcal{F}_1 \leq 1, \dots, \text{vol}_d \cap \mathcal{F}_{2d} \leq 1$ .

# Colorful versions

## Qualitative result. Puzzle:

Let finite families  $\mathcal{F}_1, \dots, \mathcal{F}_{2d}$  of convex subsets of  $\mathbb{R}^d$  be such that  $\text{vol}_d \cap \mathcal{F}_1 \leq 1, \dots, \text{vol}_d \cap \mathcal{F}_{2d} \leq 1$ .

Then there is a “rainbow” selection of sets  $F_1 \in \mathcal{F}_1, \dots, F_{2d} \in \mathcal{F}_{2d}$  such that the volume of  $F_1 \cap \dots \cap F_{2d}$  is bounded.

# Colorful versions

## Qualitative result. Puzzle:

Let finite families  $\mathcal{F}_1, \dots, \mathcal{F}_{2d}$  of convex subsets of  $\mathbb{R}^d$  be such that  $\text{vol}_d \cap \mathcal{F}_1 \leq 1, \dots, \text{vol}_d \cap \mathcal{F}_{2d} \leq 1$ .

Then there is a “rainbow” selection of sets  $F_1 \in \mathcal{F}_1, \dots, F_{2d} \in \mathcal{F}_{2d}$  such that the volume of  $F_1 \cap \dots \cap F_{2d}$  is bounded.

## Quantitative result:

We don't know the Helly-number, i.e. the number of color classes!

# Colorful versions

## Qualitative result. Puzzle:

Let finite families  $\mathcal{F}_1, \dots, \mathcal{F}_{2d}$  of convex subsets of  $\mathbb{R}^d$  be such that  $\text{vol}_d \cap \mathcal{F}_1 \leq 1, \dots, \text{vol}_d \cap \mathcal{F}_{2d} \leq 1$ .

Then there is a “rainbow” selection of sets  $F_1 \in \mathcal{F}_1, \dots, F_{2d} \in \mathcal{F}_{2d}$  such that the volume of  $F_1 \cap \dots \cap F_{2d}$  is bounded.

## Quantitative result:

We don't know the Helly-number, i.e. the number of color classes!

The best we have  $3d$  color classes (Damásdi, Földvári, Naszódi'21)!

# Log-concave functions

## Functional Bárány–Katchalski–Pach (Naszódi and I'22)

Let  $f_1, \dots, f_n$  be upper semi-continuous log-concave functions on  $\mathbb{R}^d$ . For every  $\sigma \subseteq \{1, \dots, n\}$ , let  $f_\sigma$  denote the **pointwise minimum**:

$$f_\sigma(x) = \min\{f_i(x) : i \in \sigma\}.$$

# Log-concave functions

## Functional Bárány–Katchalski–Pach (Naszódi and I'22)

Let  $f_1, \dots, f_n$  be upper semi-continuous log-concave functions on  $\mathbb{R}^d$ . For every  $\sigma \subseteq \{1, \dots, n\}$ , let  $f_\sigma$  denote the **pointwise minimum**:

$$f_\sigma(x) = \min\{f_i(x) : i \in \sigma\}.$$

Then there is a set  $\sigma \subseteq \{1, \dots, n\}$  of at most  $3d + 2$  indices such that

$$\int_{\mathbb{R}^d} f_\sigma \leq 100^d d^{2d} \int_{\mathbb{R}^d} f_{\{1, \dots, n\}}.$$

# Log-concave functions

## Functional Bárány–Katchalski–Pach (Naszódi and I'22)

Let  $f_1, \dots, f_n$  be upper semi-continuous log-concave functions on  $\mathbb{R}^d$ . For every  $\sigma \subseteq \{1, \dots, n\}$ , let  $f_\sigma$  denote the **pointwise minimum**:

$$f_\sigma(x) = \min\{f_i(x) : i \in \sigma\}.$$

Then there is a set  $\sigma \subseteq \{1, \dots, n\}$  of at most  $3d + 2$  indices such that

$$\int_{\mathbb{R}^d} f_\sigma \leq 100^d d^{2d} \int_{\mathbb{R}^d} f_{\{1, \dots, n\}}.$$

## Problem:

We don't know the Helly-number!



# Log-concave functions

## Functional Bárány–Katchalski–Pach (Naszódi and I'22)

Let  $f_1, \dots, f_n$  be upper semi-continuous log-concave functions on  $\mathbb{R}^d$ . For every  $\sigma \subseteq \{1, \dots, n\}$ , let  $f_\sigma$  denote the **pointwise minimum**:

$$f_\sigma(x) = \min\{f_i(x) : i \in \sigma\}.$$

Then there is a set  $\sigma \subseteq \{1, \dots, n\}$  of at most  $3d + 2$  indices such that

$$\int_{\mathbb{R}^d} f_\sigma \leq 100^d d^{2d} \int_{\mathbb{R}^d} f_{\{1, \dots, n\}}.$$

## Problem:

We don't know the Helly-number!

It is surely strictly less than  $3d + 2$

Thank you!