

GEOMETRIC FUNCTIONALS OF POLYCONVEX EXCURSION SETS OF POISSON SHOT NOISE PROCESSES

Vanessa Trapp

Institute of Mathematics
Hamburg University of Technology

September 29, 2023

- ▶ Excursion sets of Poisson shot noise processes
 - ▶ Definition of excursion sets of Poisson shot noise processes
 - ▶ A model with polyconvex excursion sets

- ▶ Excursion sets of Poisson shot noise processes
 - ▶ Definition of excursion sets of Poisson shot noise processes
 - ▶ A model with polyconvex excursion sets
- ▶ Geometric functionals of excursion sets
 - ▶ Asymptotic behaviour of expectation for growing observation windows
 - ▶ Asymptotic variance and central limit theorems for growing observation windows


- We consider a marked Poisson process η on $\hat{\mathbb{R}}^d := \mathbb{R}^d \times \mathbb{M}$ with intensity measure $\lambda = \gamma \lambda_d \otimes \mathbb{Q}$ for some $\gamma > 0$.

- ▶ We consider a marked Poisson process η on $\hat{\mathbb{R}}^d := \mathbb{R}^d \times \mathbb{M}$ with intensity measure $\lambda = \gamma \lambda_d \otimes \mathbb{Q}$ for some $\gamma > 0$.
- ▶ Under suitable measurability conditions, for $(g_m)_{m \in \mathbb{M}}$ with $g_m : \mathbb{R}^d \rightarrow \mathbb{R}_{\geq 0}$ a Poisson shot noise process $(f_\eta(y))_{y \in \mathbb{R}^d}$ is defined for $y \in \mathbb{R}^d$ by

$$f_\eta(y) = \sum_{(x,m) \in \eta} g_m(y - x).$$



- ▶ We consider a marked Poisson process η on $\hat{\mathbb{R}}^d := \mathbb{R}^d \times \mathbb{M}$ with intensity measure $\lambda = \gamma \lambda_d \otimes \mathbb{Q}$ for some $\gamma > 0$.
- ▶ Under suitable measurability conditions, for $(g_m)_{m \in \mathbb{M}}$ with $g_m : \mathbb{R}^d \rightarrow \mathbb{R}_{\geq 0}$ a Poisson shot noise process $(f_\eta(y))_{y \in \mathbb{R}^d}$ is defined for $y \in \mathbb{R}^d$ by

$$f_\eta(y) = \sum_{(x,m) \in \eta} g_m(y - x).$$



- ▶ We consider a marked Poisson process η on $\hat{\mathbb{R}}^d := \mathbb{R}^d \times \mathbb{M}$ with intensity measure $\lambda = \gamma \lambda_d \otimes \mathbb{Q}$ for some $\gamma > 0$.
- ▶ Under suitable measurability conditions, for $(g_m)_{m \in \mathbb{M}}$ with $g_m : \mathbb{R}^d \rightarrow \mathbb{R}_{\geq 0}$ a Poisson shot noise process $(f_\eta(y))_{y \in \mathbb{R}^d}$ is defined for $y \in \mathbb{R}^d$ by

$$f_\eta(y) = \sum_{(x,m) \in \eta} g_m(y - x).$$




- Its excursion set is given by

$$Z_u = \{y \in \mathbb{R}^d : f_\eta(y) \geq u\}.$$



- Its excursion set is given by

$$Z_u = \{y \in \mathbb{R}^d : f_\eta(y) \geq u\}.$$





- Its excursion set is given by

$$Z_u = \{y \in \mathbb{R}^d : f_\eta(y) \geq u\}.$$





- Its excursion set is given by

$$Z_u = \{y \in \mathbb{R}^d : f_\eta(y) \geq u\}.$$



- Its excursion set is given by

$$Z_u = \{y \in \mathbb{R}^d : f_\eta(y) \geq u\}.$$



- Consider a geometric functional $\varphi(Z_u \cap W)$ of the excursion set on a compact convex observation window W with $\lambda_d(W) > 0$.

- We assume that $g_m : \mathbb{R}^d \rightarrow \mathbb{R}_{\geq 0}$ is concave on its compact convex support K_m with $\lambda_d(K_m) > 0$ for $m \in \mathbb{M}$ and $m \mapsto K_m$ is measurable.

- ▶ We assume that $g_m : \mathbb{R}^d \rightarrow \mathbb{R}_{\geq 0}$ is concave on its compact convex support K_m with $\lambda_d(K_m) > 0$ for $m \in \mathbb{M}$ and $m \mapsto K_m$ is measurable.
- ▶ For $k \in \mathbb{N}$ we denote by (\mathbf{M}_k) the moment condition

$$\int_{\mathbb{M}} V_i(K_m)^k \mathbb{Q}(\mathrm{d}m) < \infty, \quad i \in \{1, \dots, d\}.$$

- ▶ We assume that $g_m : \mathbb{R}^d \rightarrow \mathbb{R}_{\geq 0}$ is concave on its compact convex support K_m with $\lambda_d(K_m) > 0$ for $m \in \mathbb{M}$ and $m \mapsto K_m$ is measurable.
- ▶ For $k \in \mathbb{N}$ we denote by (\mathbf{M}_k) the moment condition

$$\int_{\mathbb{M}} V_i(K_m)^k \mathbb{Q}(\mathrm{d}m) < \infty, \quad i \in \{1, \dots, d\}.$$

- ▶ Then, for $S_W := \{(x, m) \in \mathbb{R}^d \times \mathbb{M} : K_m + x \cap W \neq \emptyset\}$, $\eta(S_W) < \infty$ a.s.

- Let $(x_1, m_1), \dots, (x_n, m_n)$ be the points of the Poisson process in S_W . For $\hat{K}_i = K_{m_i} + x_i$, $i \in \{1, \dots, n\}$,

$$Z_u \cap W = \bigcup_{I \subseteq \{1, \dots, n\}} \left\{ y \in \bigcap_{j \in I} \hat{K}_j \cap W : \sum_{j \in I} g_{m_j}(y - x_j) \geq u \right\}.$$

- Let $(x_1, m_1), \dots, (x_n, m_n)$ be the points of the Poisson process in S_W . For $\hat{K}_i = K_{m_i} + x_i$, $i \in \{1, \dots, n\}$,

$$Z_u \cap W = \bigcup_{I \subseteq \{1, \dots, n\}} \left\{ y \in \bigcap_{j \in I} \hat{K}_j \cap W : \sum_{j \in I} g_{m_j}(y - x_j) \geq u \right\}.$$

- The excursion sets are almost surely polyconvex.

- Let $(x_1, m_1), \dots, (x_n, m_n)$ be the points of the Poisson process in S_W . For $\hat{K}_i = K_{m_i} + x_i$, $i \in \{1, \dots, n\}$,

$$Z_u \cap W = \bigcup_{I \subseteq \{1, \dots, n\}} \left\{ y \in \bigcap_{j \in I} \hat{K}_j \cap W : \sum_{j \in I} g_{m_j}(y - x_j) \geq u \right\}.$$

- The excursion sets are almost surely polyconvex.
- Boolean model arises as special case for $g_m(x) = u$ for $x \in K_m$ and $m \in \mathbb{M}$.

- We consider measurable functions $\varphi : \mathcal{R}^d \mapsto \mathbb{R}$, which are

- ▶ We consider measurable functions $\varphi : \mathcal{R}^d \mapsto \mathbb{R}$, which are
 - ▶ translation invariant, i.e. $\varphi(A + x) = \varphi(A)$ for $A \in \mathcal{R}^d, x \in \mathbb{R}^d$

- ▶ We consider measurable functions $\varphi : \mathcal{R}^d \mapsto \mathbb{R}$, which are
 - ▶ translation invariant, i.e. $\varphi(A + x) = \varphi(A)$ for $A \in \mathcal{R}^d, x \in \mathbb{R}^d$
 - ▶ additive, i.e. $\varphi(\emptyset) = 0$ and

$$\varphi(A_1 \cup A_2) = \varphi(A_1) + \varphi(A_2) - \varphi(A_1 \cap A_2)$$

for $A_1, A_2 \in \mathcal{R}^d$.

- ▶ We consider measurable functions $\varphi : \mathcal{R}^d \mapsto \mathbb{R}$, which are
 - ▶ translation invariant, i.e. $\varphi(A + x) = \varphi(A)$ for $A \in \mathcal{R}^d, x \in \mathbb{R}^d$
 - ▶ additive, i.e. $\varphi(\emptyset) = 0$ and

$$\varphi(A_1 \cup A_2) = \varphi(A_1) + \varphi(A_2) - \varphi(A_1 \cap A_2)$$

for $A_1, A_2 \in \mathcal{R}^d$.

- ▶ locally bounded, i.e. $|\varphi(A)| \leq M_\varphi$ for all $\mathcal{K}^d \ni A \subseteq [0, 1]^d$.

- ▶ We consider measurable functions $\varphi : \mathcal{R}^d \mapsto \mathbb{R}$, which are
 - ▶ translation invariant, i.e. $\varphi(A + x) = \varphi(A)$ for $A \in \mathcal{R}^d, x \in \mathbb{R}^d$
 - ▶ additive, i.e. $\varphi(\emptyset) = 0$ and

$$\varphi(A_1 \cup A_2) = \varphi(A_1) + \varphi(A_2) - \varphi(A_1 \cap A_2)$$

for $A_1, A_2 \in \mathcal{R}^d$.

- ▶ locally bounded, i.e. $|\varphi(A)| \leq M_\varphi$ for all $\mathcal{K}^d \ni A \subseteq [0, 1]^d$.
- ▶ Most prominent examples are the intrinsic volumes.

Theorem (T. 2023+)

Let $W_r = rW$ for $r \geq 1$. Under the assumption (\mathbf{M}_1) , we have

$$\lim_{r \rightarrow \infty} \frac{\mathbb{E}[\varphi(Z_u \cap W_r)]}{V_d(W_r)} = \mathbb{E}[\varphi(Z_u \cap C_0^d)],$$

where $C_0^d := [0, 1)^d$ denotes the half-open d -dimensional unit cube.

Theorem (T. 2023+)

Let $W_r = rW$ for $r \geq 1$. Under the assumption (\mathbf{M}_1) , we have

$$\lim_{r \rightarrow \infty} \frac{\mathbb{E}[\varphi(Z_u \cap W_r)]}{V_d(W_r)} = \mathbb{E}[\varphi(Z_u \cap C_0^d)],$$

where $C_0^d := [0, 1)^d$ denotes the half-open d -dimensional unit cube.

- A similar result was shown for $\mathbb{E}[\varphi(Z \cap W_r)]$, where Z is a so called standard random set in Schneider/Weil (2008).


- For a polyconvex set A let $N(A)$ denote the smallest possible number m with $A = K_1 \cup \dots \cup K_m$ for convex sets K_i , $i = 1, \dots, m$.

- ▶ For a polyconvex set A let $N(A)$ denote the smallest possible number m with $A = K_1 \cup \dots \cup K_m$ for convex sets K_i , $i = 1, \dots, m$.
- ▶ A standard random Z set satisfies $\mathbb{E}[2^{N(Z \cap [0,1]^d)}] < \infty$.

- ▶ For a polyconvex set A let $N(A)$ denote the smallest possible number m with $A = K_1 \cup \dots \cup K_m$ for convex sets K_i , $i = 1, \dots, m$.
- ▶ A standard random Z set satisfies $\mathbb{E}[2^{N(Z \cap [0,1]^d)}] < \infty$.
- ▶ $Z_u \cap W$ can be written as the number of 2^n convex sets, where $n = \eta(S_W)$, i.e. $\mathbb{E}[2^{N(Z_u \cap [0,1]^d)}]$ is not necessarily finite and therefore not a standard random set.

- Consider the example, where $g_m(x) = \frac{u}{2}$ for $x \in K_m$.

- Consider the example, where $g_m(x) = \frac{u}{2}$ for $x \in K_m$.
- In a situation as in the following picture



$N(Z_u \cap [0, 1]^d) = \frac{n^2}{4}$ and if the probability for such configurations is large enough, $\mathbb{E}[2^{N(Z_u \cap [0, 1]^d)}] = \infty$.

- For standard sets we use for $W \subset [0, 1]^d$ that $|\varphi(A)| \leq \bar{M}_\varphi$ for $A \subset W$ and hence,

$$\mathbb{E}[|\varphi(Z \cap W)|] \leq \bar{M}_\varphi \mathbb{E}[2^{N(Z \cap W)}] < \infty.$$

- For standard sets we use for $W \subset [0, 1]^d$ that $|\varphi(A)| \leq \bar{M}_\varphi$ for $A \subset W$ and hence,

$$\mathbb{E}[|\varphi(Z \cap W)|] \leq \bar{M}_\varphi \mathbb{E}[2^{N(Z \cap W)}] < \infty.$$

- For $Z_u \cap W$ we use a dynamic decomposition of W into cubes $Q_{n,z}$, which stops when $N(Z_u \cap Q_{n,z}) \leq L$ for some constant $L \in \mathbb{N}$.



- For standard sets we use for $W \subset [0, 1]^d$ that $|\varphi(A)| \leq \bar{M}_\varphi$ for $A \subset W$ and hence,

$$\mathbb{E}[|\varphi(Z \cap W)|] \leq \bar{M}_\varphi \mathbb{E}[2^{N(Z \cap W)}] < \infty.$$

- For $Z_u \cap W$ we use a dynamic decomposition of W into cubes $Q_{n,z}$, which stops when $N(Z_u \cap Q_{n,z}) \leq L$ for some constant $L \in \mathbb{N}$.



- For standard sets we use for $W \subset [0, 1]^d$ that $|\varphi(A)| \leq \bar{M}_\varphi$ for $A \subset W$ and hence,

$$\mathbb{E}[|\varphi(Z \cap W)|] \leq \bar{M}_\varphi \mathbb{E}[2^{N(Z \cap W)}] < \infty.$$

- For $Z_u \cap W$ we use a dynamic decomposition of W into cubes $Q_{n,z}$, which stops when $N(Z_u \cap Q_{n,z}) \leq L$ for some constant $L \in \mathbb{N}$.



- For standard sets we use for $W \subset [0, 1]^d$ that $|\varphi(A)| \leq \bar{M}_\varphi$ for $A \subset W$ and hence,

$$\mathbb{E}[|\varphi(Z \cap W)|] \leq \bar{M}_\varphi \mathbb{E}[2^{N(Z \cap W)}] < \infty.$$

- For $Z_u \cap W$ we use a dynamic decomposition of W into cubes $Q_{n,z}$, which stops when $N(Z_u \cap Q_{n,z}) \leq L$ for some constant $L \in \mathbb{N}$.



Theorem (T. 2023+)

Under (\mathbf{M}_2) there exists a constant $\sigma_0 \geq 0$ such that

$$\lim_{r \rightarrow \infty} \frac{\text{Var}[\varphi(Z_u \cap W_r)]}{V_d(W_r)} = \sigma_0.$$

Let N denote a standard Gaussian random variable. If $\sigma_0 > 0$ and assumption (\mathbf{M}_4) is fulfilled, there exists for $\diamond \in \{\text{Kol}, \text{Was}\}$ a constant $C > 0$ satisfying

$$d_\diamond \left(\frac{\varphi(Z_u \cap W_r) - \mathbb{E}[\varphi(Z_u \cap W_r)]}{\text{Var}[\varphi(Z_u \cap W_r)]}, N \right) \leq \frac{C}{\sqrt{V_d(W_r)}}$$

for r large enough.

- ▶ Similar results for the Boolean model in Hug/Last/Schulte (2016) or Last/Penrose (2018).

- ▶ Similar results for the Boolean model in Hug/Last/Schulte (2016) or Last/Penrose (2018).
- ▶ For specific geometric functionals as the volume of excursion sets of Poisson shot noise processes, similar results were shown in Bulinski/Spodarev/Timmermann (2012), Lachièze-Rey (2019), Lachièze-Rey/Peccati/Yang (2022), Schulte/T. (2022).

Thank you for your attention!