

# How does the sectional Poisson-Voronoi tessellation look like?


**Anna Gusakova** - Münster University  
(joint work with Zakhar Kabluchko and Christoph Thäle)

September 28, 2023

Conference on Convex Geometry and Geometric Probability,  
September 25 - 29, Salzburg


### Tessellations:

- ▶ A tessellation  $T$  in  $\mathbb{R}^d$  is a system of convex polytopes (cells), covering the space and having disjoint interiors.



## Tessellations:


- ▶ A tessellation  $T$  in  $\mathbb{R}^d$  is a system of convex polytopes (cells), covering the space and having disjoint interiors.
- ▶  $T$  is face-to-face if for any  $t_1, t_2 \in T$  we have, that  $t_1 \cap t_2$  is empty or is a face of  $t_1$  and  $t_2$ .



## Tessellations:

- ▶ A tessellation  $T$  in  $\mathbb{R}^d$  is a system of convex polytopes (cells), covering the space and having disjoint interiors.
- ▶  $T$  is face-to-face if for any  $t_1, t_2 \in T$  we have, that  $t_1 \cap t_2$  is empty or is a face of  $t_1$  and  $t_2$ .
- ▶  $\mathcal{F}_k(t)$  - set of all  $k$ -dimensional faces of a polytope  $t$ .  
Define

$$\mathcal{F}_k(T) = \bigcup_{t \in T} \mathcal{F}_k(t), 0 \leq k \leq d.$$




## Tessellations:

- ▶ A tessellation  $T$  in  $\mathbb{R}^d$  is a system of convex polytopes (cells), covering the space and having disjoint interiors.
- ▶  $T$  is face-to-face if for any  $t_1, t_2 \in T$  we have, that  $t_1 \cap t_2$  is empty or is a face of  $t_1$  and  $t_2$ .
- ▶  $\mathcal{F}_k(t)$  - set of all  $k$ -dimensional faces of a polytope  $t$ .  
Define

$$\mathcal{F}_k(T) = \bigcup_{t \in T} \mathcal{F}_k(t), 0 \leq k \leq d.$$

- ▶  $T$  is called normal if any  $F \in \mathcal{F}_k(T)$  is contained in exactly  $d - k + 1$  cells.




## Tessellations:

- ▶ A tessellation  $T$  in  $\mathbb{R}^d$  is a system of convex polytopes (cells), covering the space and having disjoint interiors.
- ▶  $T$  is face-to-face if for any  $t_1, t_2 \in T$  we have, that  $t_1 \cap t_2$  is empty or is a face of  $t_1$  and  $t_2$ .
- ▶  $\mathcal{F}_k(t)$  - set of all  $k$ -dimensional faces of a polytope  $t$ .  
Define

$$\mathcal{F}_k(T) = \bigcup_{t \in T} \mathcal{F}_k(t), 0 \leq k \leq d.$$


- ▶  $T$  is called normal if any  $F \in \mathcal{F}_k(T)$  is contained in exactly  $d - k + 1$  cells.
- ▶  $T$  is called simplicial if all its cells are simplices.



## Tessellations:

- ▶ A tessellation  $T$  in  $\mathbb{R}^d$  is a system of convex polytopes (cells), covering the space and having disjoint interiors.
- ▶  $T$  is face-to-face if for any  $t_1, t_2 \in T$  we have, that  $t_1 \cap t_2$  is empty or is a face of  $t_1$  and  $t_2$ .
- ▶  $\mathcal{F}_k(t)$  - set of all  $k$ -dimensional faces of a polytope  $t$ .  
Define

$$\mathcal{F}_k(T) = \bigcup_{t \in T} \mathcal{F}_k(t), 0 \leq k \leq d.$$



- ▶  $T$  is called normal if any  $F \in \mathcal{F}_k(T)$  is contained in exactly  $d - k + 1$  cells.
- ▶  $T$  is called simplicial if all its cells are simplices.

**Poisson point process**  $\eta$  in  $\mathbb{R}^d$  with intensity measure  $\mu$ :

- ▶  $\forall A \in \mathcal{B}(\mathbb{R}^d), \eta(A) \sim \text{Poi}(\mu(A));$
- ▶  $\forall$  mutually disjoint subsets  $A_1, \dots, A_m \in \mathcal{B}(\mathbb{R}^d), \eta(A_1), \dots, \eta(A_m)$  are independent.

# Poisson-Voronoi tessellation: construction

Let  $\eta$  be a PPP in  $\mathbb{R}^d$  with intensity measure  $\gamma \text{Leb}_d$ ,  $\gamma > 0$ .





# Poisson-Voronoi tessellation: construction

Let  $\eta$  be a PPP in  $\mathbb{R}^d$  with intensity measure  $\gamma \text{Leb}_d$ ,  $\gamma > 0$ .

**Voronoi cell** of (nuclei)  $v \in \eta$ :  $V(v, \eta) := \{z \in \mathbb{R}^d : \|z - v\|^2 \leq \|z - v'\|^2 \text{ for all } v' \in \eta\}$ .



Let  $\eta$  be a PPP in  $\mathbb{R}^d$  with intensity measure  $\gamma \text{Leb}_d$ ,  $\gamma > 0$ .

**Voronoi cell** of (nuclei)  $v \in \eta$ :  $V(v, \eta) := \{z \in \mathbb{R}^d : \|z - v\|^2 \leq \|z - v'\|^2 \text{ for all } v' \in \eta\}$ .

Let  $\eta$  be a PPP in  $\mathbb{R}^d$  with intensity measure  $\gamma \text{Leb}_d$ ,  $\gamma > 0$ .

**Voronoi cell** of (nuclei)  $v \in \eta$ :  $V(v, \eta) := \{z \in \mathbb{R}^d : \|z - v\|^2 \leq \|z - v'\|^2 \text{ for all } v' \in \eta\}$ .

**Poisson-Voronoi tessellation**:  $\mathcal{V}_\gamma^d := \{V(v, \eta) : v \in \eta\}$ .

Let  $\eta$  be a PPP in  $\mathbb{R}^d$  with intensity measure  $\gamma \text{Leb}_d$ ,  $\gamma > 0$ .


**Voronoi cell** of (nuclei)  $v \in \eta$ :  $V(v, \eta) := \{z \in \mathbb{R}^d : \|z - v\|^2 \leq \|z - v'\|^2 \text{ for all } v' \in \eta\}$ .

**Poisson-Voronoi tessellation**:  $\mathcal{V}_\gamma^d := \{V(v, \eta) : v \in \eta\}$ .

Fact:  $\mathcal{V}_\gamma^d$  is almost surely face-to-face, normal random tessellation and  $\mathcal{V}_\gamma^d$  is stationary.

# Characteristics of stationary random tessellation

- ▶  $\mathcal{T}$  is a stationary random tessellation.




# Characteristics of stationary random tessellation

- ▶  $\mathcal{T}$  is a stationary random tessellation.
- ▶  **$k$ -cell intensity of  $\mathcal{T}$ :**

$$\gamma_k(\mathcal{T}) = \mathbb{E} \sum_{F \in \mathcal{F}_k(\mathcal{T})} \mathbf{1}\{z(F, \mathcal{T}) \in [0, 1]^d\},$$

where  $z : \{\text{polytopes}\} \times \{\text{tessellations}\} \mapsto \mathbb{R}^d$  be s.t.  
 $z(t + x, \mathcal{T} + x) = z(t, \mathcal{T}) + x$  for all  $x \in \mathbb{R}^d$ .




# Characteristics of stationary random tessellation

- ▶  $\mathcal{T}$  is a stationary random tessellation.
- ▶  **$k$ -cell intensity of  $\mathcal{T}$ :**

$$\gamma_k(\mathcal{T}) = \mathbb{E} \sum_{F \in \mathcal{F}_k(\mathcal{T})} \mathbf{1}\{z(F, \mathcal{T}) \in [0, 1]^d\},$$

where  $z : \{\text{polytopes}\} \times \{\text{tessellations}\} \mapsto \mathbb{R}^d$  be s.t.  
 $z(t + x, \mathcal{T} + x) = z(t, \mathcal{T}) + x$  for all  $x \in \mathbb{R}^d$ .




# Characteristics of stationary random tessellation

- ▶  $\mathcal{T}$  is a stationary random tessellation.
- ▶  **$k$ -cell intensity of  $\mathcal{T}$ :**

$$\gamma_k(\mathcal{T}) = \mathbb{E} \sum_{F \in \mathcal{F}_k(\mathcal{T})} \mathbf{1}\{z(F, \mathcal{T}) \in [0, 1]^d\},$$

where  $z : \{\text{polytopes}\} \times \{\text{tessellations}\} \mapsto \mathbb{R}^d$  be s.t.  
 $z(t + x, \mathcal{T} + x) = z(t, \mathcal{T}) + x$  for all  $x \in \mathbb{R}^d$ .



$k = 2$




# Characteristics of stationary random tessellation

- ▶  $\mathcal{T}$  is a stationary random tessellation.
- ▶  **$k$ -cell intensity of  $\mathcal{T}$ :**

$$\gamma_k(\mathcal{T}) = \mathbb{E} \sum_{F \in \mathcal{F}_k(\mathcal{T})} \mathbf{1}\{z(F, \mathcal{T}) \in [0, 1]^d\},$$

where  $z : \{\text{polytopes}\} \times \{\text{tessellations}\} \mapsto \mathbb{R}^d$  be s.t.  
 $z(t + x, \mathcal{T} + x) = z(t, \mathcal{T}) + x$  for all  $x \in \mathbb{R}^d$ .




# Characteristics of stationary random tessellation

- ▶  $\mathcal{T}$  is a stationary random tessellation.
- ▶  **$k$ -cell intensity of  $\mathcal{T}$ :**

$$\gamma_k(\mathcal{T}) = \mathbb{E} \sum_{F \in \mathcal{F}_k(\mathcal{T})} \mathbf{1}\{z(F, \mathcal{T}) \in [0, 1]^d\},$$

where  $z : \{\text{polytopes}\} \times \{\text{tessellations}\} \mapsto \mathbb{R}^d$  be s.t.  
 $z(t + x, \mathcal{T} + x) = z(t, \mathcal{T}) + x$  for all  $x \in \mathbb{R}^d$ .



- ▶ **Typical cell of  $\mathcal{T}$ :**  $Z(\mathcal{T})$  is a random polytope with distribution

$$\mathbb{P}_{\mathcal{T}}^{\text{typ}}(\cdot) = \frac{1}{\gamma_d(\mathcal{T})} \mathbb{E} \sum_{t \in \mathcal{T} : z(t, \mathcal{T}) \in [0, 1]^d} \mathbf{1}\{t - z(t, \mathcal{T}) \in \cdot\}.$$

For  $\mathcal{V}_{\gamma}^d$  we choose  $z(V(v, \eta), \mathcal{V}_{\gamma}^d) = v$ .

# Characteristics of stationary random tessellation

- ▶  $\mathcal{T}$  is a stationary random tessellation.
- ▶  **$k$ -cell intensity of  $\mathcal{T}$ :**


$$\gamma_k(\mathcal{T}) = \mathbb{E} \sum_{F \in \mathcal{F}_k(\mathcal{T})} \mathbf{1}\{z(F, \mathcal{T}) \in [0, 1]^d\},$$

where  $z : \{\text{polytopes}\} \times \{\text{tessellations}\} \mapsto \mathbb{R}^d$  be s.t.  
 $z(t + x, \mathcal{T} + x) = z(t, \mathcal{T}) + x$  for all  $x \in \mathbb{R}^d$ .

- ▶ **Typical cell of  $\mathcal{T}$ :**  $Z(\mathcal{T})$  is a random polytope with distribution

$$\mathbb{P}_{\mathcal{T}}^{\text{typ}}(\cdot) = \frac{1}{\gamma_d(\mathcal{T})} \mathbb{E} \sum_{t \in \mathcal{T} : z(t, \mathcal{T}) \in [0, 1]^d} \mathbf{1}\{t - z(t, \mathcal{T}) \in \cdot\}.$$

For  $\mathcal{V}_{\gamma}^d$  we choose  $z(V(v, \eta), \mathcal{V}_{\gamma}^d) = v$ .




# Characteristics of stationary random tessellation

- ▶  $\mathcal{T}$  is a stationary random tessellation.

- ▶  **$k$ -cell intensity of  $\mathcal{T}$ :**

$$\gamma_k(\mathcal{T}) = \mathbb{E} \sum_{F \in \mathcal{F}_k(\mathcal{T})} \mathbf{1}\{z(F, \mathcal{T}) \in [0, 1]^d\},$$

where  $z : \{\text{polytopes}\} \times \{\text{tessellations}\} \mapsto \mathbb{R}^d$  be s.t.  
 $z(t + x, \mathcal{T} + x) = z(t, \mathcal{T}) + x$  for all  $x \in \mathbb{R}^d$ .



- ▶ **Typical cell of  $\mathcal{T}$ :**  $Z(\mathcal{T})$  is a random polytope with distribution

$$\mathbb{P}_{\mathcal{T}}^{\text{typ}}(\cdot) = \frac{1}{\gamma_d(\mathcal{T})} \mathbb{E} \sum_{t \in \mathcal{T} : z(t, \mathcal{T}) \in [0, 1]^d} \mathbf{1}\{t - z(t, \mathcal{T}) \in \cdot\}.$$

For  $\mathcal{V}_{\gamma}^d$  we choose  $z(V(v, \eta), \mathcal{V}_{\gamma}^d) = v$ .

- ▶ Functions of interest:

- ▶  $f_k(Z(\mathcal{T}))$  – number of  $k$ -dimensional faces;


# Characteristics of stationary random tessellation

- ▶  $\mathcal{T}$  is a stationary random tessellation.

- ▶  **$k$ -cell intensity of  $\mathcal{T}$ :**

$$\gamma_k(\mathcal{T}) = \mathbb{E} \sum_{F \in \mathcal{F}_k(\mathcal{T})} \mathbf{1}\{z(F, \mathcal{T}) \in [0, 1]^d\},$$

where  $z : \{\text{polytopes}\} \times \{\text{tessellations}\} \mapsto \mathbb{R}^d$  be s.t.  
 $z(t + x, \mathcal{T} + x) = z(t, \mathcal{T}) + x$  for all  $x \in \mathbb{R}^d$ .



- ▶ **Typical cell of  $\mathcal{T}$ :**  $Z(\mathcal{T})$  is a random polytope with distribution

$$\mathbb{P}_{\mathcal{T}}^{\text{typ}}(\cdot) = \frac{1}{\gamma_d(\mathcal{T})} \mathbb{E} \sum_{t \in \mathcal{T} : z(t, \mathcal{T}) \in [0, 1]^d} \mathbf{1}\{t - z(t, \mathcal{T}) \in \cdot\}.$$

For  $\mathcal{V}_{\gamma}^d$  we choose  $z(V(v, \eta), \mathcal{V}_{\gamma}^d) = v$ .

- ▶ Functions of interest:
  - ▶  $f_k(Z(\mathcal{T}))$  – number of  $k$ -dimensional faces;
  - ▶  $V_j(Z(\mathcal{T}))$  –  $j$ -th intrinsic volume.


# Poisson-Voronoi tessellation $\mathcal{V}_\gamma^d$ : overview

Use simplified notation:  $Z_d := Z(\mathcal{V}_\gamma^d)$ ,  $\gamma_k := \gamma_k(\mathcal{V}_\gamma^d)$ .

# Poisson-Voronoi tessellation $\mathcal{V}_\gamma^d$ : overview

Use simplified notation:  $Z_d := Z(\mathcal{V}_\gamma^d)$ ,  $\gamma_k := \gamma_k(\mathcal{V}_\gamma^d)$ .


- Trivial cases:  $\gamma_d = \gamma$ ,  $\mathbb{E}V_d(Z_d) = \gamma^{-1}$ .



# Poisson-Voronoi tessellation $\mathcal{V}_\gamma^d$ : overview

Use simplified notation:  $Z_d := Z(\mathcal{V}_\gamma^d)$ ,  $\gamma_k := \gamma_k(\mathcal{V}_\gamma^d)$ .

- ▶ Trivial cases:  $\gamma_d = \gamma$ ,  $\mathbb{E}V_d(Z_d) = \gamma^{-1}$ .
- ▶  $\mathcal{V}_\gamma^d$  is normal:  $(d - k + 1)\gamma_k = \gamma \mathbb{E}f_k(Z_d)$  and  $2\gamma_1 = (d + 1)\gamma_0$ .






# Poisson-Voronoi tessellation $\mathcal{V}_\gamma^d$ : overview

Use simplified notation:  $Z_d := Z(\mathcal{V}_\gamma^d)$ ,  $\gamma_k := \gamma_k(\mathcal{V}_\gamma^d)$ .

- ▶ Trivial cases:  $\gamma_d = \gamma$ ,  $\mathbb{E}V_d(Z_d) = \gamma^{-1}$ .
- ▶  $\mathcal{V}_\gamma^d$  is normal:  $(d - k + 1)\gamma_k = \gamma \mathbb{E}f_k(Z_d)$  and  $2\gamma_1 = (d + 1)\gamma_0$ .
- ▶ Gilbert, 1962; Miles, 1970; Møller, 1989:

$$\mathbb{E}f_0(Z_d) = 2^d \pi^{\frac{d-1}{2}} \frac{\Gamma(\frac{d^2}{2} + \frac{1}{2})}{\Gamma(\frac{d^2}{2} + 1)} \frac{\Gamma(\frac{d}{2} + 1)^d}{\Gamma(\frac{d+1}{2})^d}$$

$$\mathbb{E}V_{d-1}(Z_d) = 2^{d-1} \frac{\Gamma(\frac{d}{2})\Gamma(2 - \frac{1}{d})}{\Gamma(d - \frac{1}{2})} \Gamma(\frac{d}{2} + 1)^{1 - \frac{1}{d}} \gamma^{-1 + \frac{1}{d}}.$$




# Poisson-Voronoi tessellation $\mathcal{V}_\gamma^d$ : overview

Use simplified notation:  $Z_d := Z(\mathcal{V}_\gamma^d)$ ,  $\gamma_k := \gamma_k(\mathcal{V}_\gamma^d)$ .

- ▶ Trivial cases:  $\gamma_d = \gamma$ ,  $\mathbb{E}V_d(Z_d) = \gamma^{-1}$ .
- ▶  $\mathcal{V}_\gamma^d$  is normal:  $(d - k + 1)\gamma_k = \gamma \mathbb{E}f_k(Z_d)$  and  $2\gamma_1 = (d + 1)\gamma_0$ .
- ▶ Gilbert, 1962; Miles, 1970; Møller, 1989:

$$\mathbb{E}f_0(Z_d) = C_1(d)$$


$$\mathbb{E}V_{d-1}(Z_d) = C_2(d)\gamma^{-1+\frac{1}{d}}.$$



# Poisson-Voronoi tessellation $\mathcal{V}_\gamma^d$ : overview

Use simplified notation:  $Z_d := Z(\mathcal{V}_\gamma^d)$ ,  $\gamma_k := \gamma_k(\mathcal{V}_\gamma^d)$ .

- ▶ Trivial cases:  $\gamma_d = \gamma$ ,  $\mathbb{E}V_d(Z_d) = \gamma^{-1}$ .
- ▶  $\mathcal{V}_\gamma^d$  is normal:  $(d - k + 1)\gamma_k = \gamma \mathbb{E}f_k(Z_d)$  and  $2\gamma_1 = (d + 1)\gamma_0$ .
- ▶ Gilbert, 1962; Miles, 1970; Møller, 1989:




$$\begin{aligned}\mathbb{E}f_0(Z_d) &= C_1(d) & \Rightarrow & \gamma_0, \gamma_1, \mathbb{E}f_1(Z_d), \\ \mathbb{E}V_{d-1}(Z_d) &= C_2(d)\gamma^{-1+\frac{1}{d}}.\end{aligned}$$

# Poisson-Voronoi tessellation $\mathcal{V}_\gamma^d$ : overview

Use simplified notation:  $Z_d := Z(\mathcal{V}_\gamma^d)$ ,  $\gamma_k := \gamma_k(\mathcal{V}_\gamma^d)$ .

- ▶ Trivial cases:  $\gamma_d = \gamma$ ,  $\mathbb{E}V_d(Z_d) = \gamma^{-1}$ .
- ▶  $\mathcal{V}_\gamma^d$  is normal:  $(d - k + 1)\gamma_k = \gamma \mathbb{E}f_k(Z_d)$  and  $2\gamma_1 = (d + 1)\gamma_0$ .
- ▶ Gilbert, 1962; Miles, 1970; Møller, 1989:



$$\begin{aligned}\mathbb{E}f_0(Z_d) &= C_1(d) & \Rightarrow & \gamma_0, \gamma_1, \mathbb{E}f_1(Z_d), \\ \mathbb{E}V_{d-1}(Z_d) &= C_2(d)\gamma^{-1+\frac{1}{d}}.\end{aligned}$$


Define the function:

$$J_{n,k}\left(\beta + \frac{1}{2}\right) = c_{n(n+2\beta)} \int_{-\infty}^{\infty} (\cosh u)^{-n(n+2\beta)-2} \left[ \frac{1}{2} + i \int_0^u c_{n+2\beta-1} (\cosh v)^{n+2\beta} dv \right]^{n-k} du,$$

where  $c_\alpha = \pi^{-\frac{1}{2}} \Gamma\left(\frac{\alpha+3}{2}\right) \Gamma\left(\frac{\alpha+2}{2}\right)^{-1}$ .

Use simplified notation:  $Z_d := Z(\mathcal{V}_\gamma^d)$ ,  $\gamma_k := \gamma_k(\mathcal{V}_\gamma^d)$ .

- ▶ Trivial cases:  $\gamma_d = \gamma$ ,  $\mathbb{E}V_d(Z_d) = \gamma^{-1}$ .
- ▶  $\mathcal{V}_\gamma^d$  is normal:  $(d - k + 1)\gamma_k = \gamma \mathbb{E}f_k(Z_d)$  and  $2\gamma_1 = (d + 1)\gamma_0$ .
- ▶ Gilbert, 1962; Miles, 1970; Møller, 1989:



$$\begin{aligned} \mathbb{E}f_0(Z_d) &= C_1(d) & \Rightarrow & \gamma_0, \gamma_1, \mathbb{E}f_1(Z_d), \\ \mathbb{E}V_{d-1}(Z_d) &= C_2(d)\gamma^{-1+\frac{1}{d}}. \end{aligned}$$

Define the function:

$$J_{n,k}\left(\beta + \frac{1}{2}\right) = c_{n(n+2\beta)} \int_{-\infty}^{\infty} (\cosh u)^{-n(n+2\beta)-2} \left[ \frac{1}{2} + i \int_0^u c_{n+2\beta-1} (\cosh v)^{n+2\beta} dv \right]^{n-k} du,$$

where  $c_\alpha = \pi^{-\frac{1}{2}} \Gamma\left(\frac{\alpha+3}{2}\right) \Gamma\left(\frac{\alpha+2}{2}\right)^{-1}$ .


- ▶ Kabluchko, Thäle, Zaporozhets, 2020; Kabluchko, 2021:

$$\mathbb{E}f_k(Z_d) = C_1(d) \binom{d}{k} J_{d+1, d-k+1}\left(-\frac{1}{2}\right)$$

# Poisson-Voronoi tessellation $\mathcal{V}_\gamma^d$ : overview

Use simplified notation:  $Z_d := Z(\mathcal{V}_\gamma^d)$ ,  $\gamma_k := \gamma_k(\mathcal{V}_\gamma^d)$ .

- ▶ Trivial cases:  $\gamma_d = \gamma$ ,  $\mathbb{E}V_d(Z_d) = \gamma^{-1}$ .
- ▶  $\mathcal{V}_\gamma^d$  is normal:  $(d - k + 1)\gamma_k = \gamma \mathbb{E}f_k(Z_d)$  and  $2\gamma_1 = (d + 1)\gamma_0$ .
- ▶ Gilbert, 1962; Miles, 1970; Møller, 1989:



$$\begin{aligned}\mathbb{E}f_0(Z_d) &= C_1(d) & \Rightarrow & \gamma_0, \gamma_1, \mathbb{E}f_1(Z_d), \\ \mathbb{E}V_{d-1}(Z_d) &= C_2(d)\gamma^{-1+\frac{1}{d}}.\end{aligned}$$

Define the function:

$$J_{n,k}\left(\beta + \frac{1}{2}\right) = c_{n(n+2\beta)} \int_{-\infty}^{\infty} (\cosh u)^{-n(n+2\beta)-2} \left[ \frac{1}{2} + i \int_0^u c_{n+2\beta-1} (\cosh v)^{n+2\beta} dv \right]^{n-k} du,$$

where  $c_\alpha = \pi^{-\frac{1}{2}} \Gamma\left(\frac{\alpha+3}{2}\right) \Gamma\left(\frac{\alpha+2}{2}\right)^{-1}$ .

- ▶ Kabluchko, Thäle, Zaporozhets, 2020; Kabluchko, 2021:

$$\mathbb{E}f_k(Z_d) = C_1(d) \binom{d}{k} J_{d+1, d-k+1}\left(-\frac{1}{2}\right) \Rightarrow \gamma_k.$$

- Kabluchko, Thäle, Zaporozhets, 2020; Kabluchko, 2021:

$$\mathbb{E}f_k(Z_d) = C_1(d) \binom{d}{k} J_{d+1, d-k+1} \left( -\frac{1}{2} \right).$$

Idea of the proof:

- Kabluchko, Thäle, Zaporozhets, 2020; Kabluchko, 2021:


$$\mathbb{E}f_k(Z_d) = C_1(d) \binom{d}{k} J_{d+1, d-k+1} \left( -\frac{1}{2} \right).$$

Idea of the proof:

- Suitable stochastic representation of  $Z_d$

$$(Z_d)^\circ \stackrel{d}{=} \text{conv}(\Pi_{d,d})$$

where  $\Pi_{d,d}$  is a PPP with intensity measure  $\|x\|^{-d-1}dx$ .





- Kabluchko, Thäle, Zaporozhets, 2020; Kabluchko, 2021:


$$\mathbb{E}f_k(Z_d) = C_1(d) \binom{d}{k} J_{d+1, d-k+1} \left( -\frac{1}{2} \right).$$

Idea of the proof:

- Suitable stochastic representation of  $Z_d$

$$(Z_d)^\circ \stackrel{d}{=} \text{conv}(\Pi_{d,d}) \quad \Rightarrow \quad f_k(Z_d) = f_{d-1-k}(\text{conv}(\Pi_{d,d})),$$

where  $\Pi_{d,d}$  is a PPP with intensity measure  $\|x\|^{-d-1}dx$ .



- Kabluchko, Thäle, Zaporozhets, 2020; Kabluchko, 2021:

$$\mathbb{E}f_k(Z_d) = C_1(d) \binom{d}{k} J_{d+1, d-k+1} \left( -\frac{1}{2} \right).$$

Idea of the proof:

- Suitable stochastic representation of  $Z_d$


$$(Z_d)^\circ \stackrel{d}{=} \text{conv}(\Pi_{d,d}) \quad \Rightarrow \quad f_k(Z_d) = f_{d-1-k}(\text{conv}(\Pi_{d,d})),$$

where  $\Pi_{d,d}$  is a PPP with intensity measure  $\|x\|^{-d-1}dx$ .

- Let  $X_1, \dots, X_n$  be i.i.d. points in  $\mathbb{R}^d$  with density function

$$x \mapsto \pi^{-\frac{d}{2}} \Gamma(d) \Gamma\left(\frac{d}{2}\right)^{-1} (1 + \|x\|^2)^{-d},$$

and denote  $P_{n,d} := \text{conv}(X_1, \dots, X_n) - \beta'$ -polytope.



- Kabluchko, Thäle, Zaporozhets, 2020; Kabluchko, 2021:

$$\mathbb{E}f_k(Z_d) = C_1(d) \binom{d}{k} J_{d+1, d-k+1} \left( -\frac{1}{2} \right).$$

Idea of the proof:

- Suitable stochastic representation of  $Z_d$

$$(Z_d)^\circ \stackrel{d}{=} \text{conv}(\Pi_{d,d}) \quad \Rightarrow \quad f_k(Z_d) = f_{d-1-k}(\text{conv}(\Pi_{d,d})),$$


where  $\Pi_{d,d}$  is a PPP with intensity measure  $\|x\|^{-d-1}dx$ .

- Let  $X_1, \dots, X_n$  be i.i.d. points in  $\mathbb{R}^d$  with density function

$$x \mapsto \pi^{-\frac{d}{2}} \Gamma(d) \Gamma\left(\frac{d}{2}\right)^{-1} (1 + \|x\|^2)^{-d},$$

and denote  $P_{n,d} := \text{conv}(X_1, \dots, X_n) - \beta'$ -polytope.

- $\mathbb{E}f_{d-1-k}(\text{conv}(\Pi_{d,d})) = \lim_{n \rightarrow \infty} \mathbb{E}f_{d-1-k}(P_{n,d})$



- Kabluchko, Thäle, Zaporozhets, 2020; Kabluchko, 2021:

$$\mathbb{E}f_k(Z_d) = C_1(d) \binom{d}{k} J_{d+1, d-k+1} \left( -\frac{1}{2} \right).$$

Idea of the proof:

- Suitable stochastic representation of  $Z_d$

$$(Z_d)^\circ \stackrel{d}{=} \text{conv}(\Pi_{d,d}) \quad \Rightarrow \quad f_k(Z_d) = f_{d-1-k}(\text{conv}(\Pi_{d,d})),$$


where  $\Pi_{d,d}$  is a PPP with intensity measure  $\|x\|^{-d-1}dx$ .

- Let  $X_1, \dots, X_n$  be i.i.d. points in  $\mathbb{R}^d$  with density function

$$x \mapsto \pi^{-\frac{d}{2}} \Gamma(d) \Gamma\left(\frac{d}{2}\right)^{-1} (1 + \|x\|^2)^{-d},$$

and denote  $P_{n,d} := \text{conv}(X_1, \dots, X_n) - \beta'$ -polytope.

- $\mathbb{E}f_{d-1-k}(\text{conv}(\Pi_{d,d})) = \lim_{n \rightarrow \infty} \mathbb{E}f_{d-1-k}(P_{n,d}) + \text{exact formulas for } \mathbb{E}f_{d-1-k}(P_{n,d}).$



- Kabluchko, Thäle, Zaporozhets, 2020; Kabluchko, 2021:

$$\mathbb{E}f_k(Z_d) = C_1(d) \binom{d}{k} J_{d+1, d-k+1} \left( -\frac{1}{2} \right).$$

Idea of the proof:

- Suitable stochastic representation of  $Z_d$

$$(Z_d)^\circ \stackrel{d}{=} \text{conv}(\Pi_{d,d}) \quad \Rightarrow \quad f_k(Z_d) = f_{d-1-k}(\text{conv}(\Pi_{d,d})),$$


where  $\Pi_{d,d}$  is a PPP with intensity measure  $\|x\|^{-d-1}dx$ .

- Let  $X_1, \dots, X_n$  be i.i.d. points in  $\mathbb{R}^d$  with density function

$$x \mapsto \pi^{-\frac{d}{2}} \Gamma(d) \Gamma\left(\frac{d}{2}\right)^{-1} (1 + \|x\|^2)^{-d},$$

and denote  $P_{n,d} := \text{conv}(X_1, \dots, X_n) - \beta'$ -polytope.

- $\mathbb{E}f_{d-1-k}(\text{conv}(\Pi_{d,d})) = \lim_{n \rightarrow \infty} \mathbb{E}f_{d-1-k}(P_{n,d}) + \text{exact formulas for } \mathbb{E}f_{d-1-k}(P_{n,d})$ .



This approach doesn't help to study expected intrinsic volumes of the typical cell.

Given a  $\ell$ -dimensional affine subspace  $L_\ell \subset \mathbb{R}^d$  define the intersection tessellation

$$\mathcal{V}_\gamma^d \cap L_\ell := \{t \cap L_\ell : t \in \mathcal{V}_\gamma^d\}.$$

Given a  $\ell$ -dimensional affine subspace  $L_\ell \subset \mathbb{R}^d$  define the intersection tessellation

$$\mathcal{V}_\gamma^d \cap L_\ell := \{t \cap L_\ell : t \in \mathcal{V}_\gamma^d\}.$$

Useful formula:  $\mathbb{E}V_{d-\ell}(Z_d) = \frac{d!\Gamma(\frac{\ell}{2}+1)\Gamma(\frac{d-\ell}{2}+1)}{\ell!(d-\ell)!\Gamma(\frac{d}{2}+1)}\gamma_\ell(\mathcal{V}_\gamma^d \cap L_\ell)\gamma^{-1}.$

Given a  $\ell$ -dimensional affine subspace  $L_\ell \subset \mathbb{R}^d$  define the intersection tessellation

$$\mathcal{V}_\gamma^d \cap L_\ell := \{t \cap L_\ell : t \in \mathcal{V}_\gamma^d\}.$$

Useful formula:  $\mathbb{E}V_{d-\ell}(Z_d) = C_3(d, \ell)\gamma_\ell(\mathcal{V}_\gamma^d \cap L_\ell)\gamma^{-1}$ .



Given a  $\ell$ -dimensional affine subspace  $L_\ell \subset \mathbb{R}^d$  define the intersection tessellation

$$\mathcal{V}_\gamma^d \cap L_\ell := \{t \cap L_\ell : t \in \mathcal{V}_\gamma^d\}.$$

Useful formula:  $\mathbb{E}V_{d-\ell}(Z_d) = C_3(d, \ell) \gamma_\ell(\mathcal{V}_\gamma^d \cap L_\ell) \gamma^{-1}$ .

Intuitional proof: Crofton's formula  $\mathbb{E}V_{d-\ell}(Z_d) = C_3(d, \ell) \int_{A(d, \ell)} \mathbb{E}\chi(Z_d \cap L) \mu_\ell(dL)$ .

Given a  $\ell$ -dimensional affine subspace  $L_\ell \subset \mathbb{R}^d$  define the intersection tessellation

$$\mathcal{V}_\gamma^d \cap L_\ell := \{t \cap L_\ell : t \in \mathcal{V}_\gamma^d\}.$$

Useful formula:  $\mathbb{E}V_{d-\ell}(Z_d) = C_3(d, \ell)\gamma_\ell(\mathcal{V}_\gamma^d \cap L_\ell)\gamma^{-1}$ .

Intuitional proof: Crofton's formula  $\mathbb{E}V_{d-\ell}(Z_d) = C_3(d, \ell) \int_{A(d, \ell)} \mathbb{E} \mathbf{1}\{Z_d \cap L \neq \emptyset\} \mu_\ell(dL)$ .


# Sectional Poisson-Voronoi tessellation

Given a  $\ell$ -dimensional affine subspace  $L_\ell \subset \mathbb{R}^d$  define the intersection tessellation

$$\mathcal{V}_\gamma^d \cap L_\ell := \{t \cap L_\ell : t \in \mathcal{V}_\gamma^d\}.$$

Useful formula:  $\mathbb{E}V_{d-\ell}(Z_d) = C_3(d, \ell) \gamma_\ell(\mathcal{V}_\gamma^d \cap L_\ell) \gamma^{-1}$ .

Intuitional proof: Crofton's formula  $\mathbb{E}V_{d-\ell}(Z_d) = C_3(d, \ell) \int_{A(d, \ell)} \mathbb{E} \mathbf{1}\{Z_d \cap L \neq \emptyset\} \mu_\ell(dL)$ .




## Sectional Poisson-Voronoi tessellation

Given a  $\ell$ -dimensional affine subspace  $L_\ell \subset \mathbb{R}^d$  define the intersection tessellation

$$\mathcal{V}_\gamma^d \cap L_\ell := \{t \cap L_\ell : t \in \mathcal{V}_\gamma^d\}.$$

Useful formula:  $\mathbb{E}V_{d-\ell}(Z_d) = C_3(d, \ell)\gamma_\ell(\mathcal{V}_\gamma^d \cap L_\ell)\gamma^{-1}$ .

Intuitional proof: Crofton's formula  $\mathbb{E}V_{d-\ell}(Z_d) = C_3(d, \ell)\gamma_\ell(\mathcal{V}_\gamma^d \cap L_\ell)\gamma^{-1}$ .



## Sectional Poisson-Voronoi tessellation

Given a  $\ell$ -dimensional affine subspace  $L_\ell \subset \mathbb{R}^d$  define the intersection tessellation

$$\mathcal{V}_\gamma^d \cap L_\ell := \{t \cap L_\ell : t \in \mathcal{V}_\gamma^d\}.$$

Useful formula:  $\mathbb{E}V_{d-\ell}(Z_d) = C_3(d, \ell)\gamma_\ell(\mathcal{V}_\gamma^d \cap L_\ell)\gamma^{-1}$ .

**Question:** Is the sectional Poisson-Voronoi tessellation  $\mathcal{V}_\gamma^d \cap L_\ell$  a Voronoi tessellation?


# Sectional Poisson-Voronoi tessellation

Given a  $\ell$ -dimensional affine subspace  $L_\ell \subset \mathbb{R}^d$  define the intersection tessellation


$$\mathcal{V}_\gamma^d \cap L_\ell := \{t \cap L_\ell : t \in \mathcal{V}_\gamma^d\}.$$

Useful formula:  $\mathbb{E}V_{d-\ell}(Z_d) = C_3(d, \ell)\gamma_\ell(\mathcal{V}_\gamma^d \cap L_\ell)\gamma^{-1}$ .

**Question:** Is the sectional Poisson-Voronoi tessellation  $\mathcal{V}_\gamma^d \cap L_\ell$  a Voronoi tessellation?



3D Poisson-Voronoi tessellation



2D Poisson-Voronoi tessellation

## Sectional Poisson-Voronoi tessellation

Given a  $\ell$ -dimensional affine subspace  $L_\ell \subset \mathbb{R}^d$  define the intersection tessellation

$$\mathcal{V}_\gamma^d \cap L_\ell := \{t \cap L_\ell : t \in \mathcal{V}_\gamma^d\}.$$

Useful formula:  $\mathbb{E}V_{d-\ell}(Z_d) = C_3(d, \ell) \gamma_\ell (\mathcal{V}_\gamma^d \cap L_\ell) \gamma^{-1}$ .

**Question:** Is the sectional Poisson-Voronoi tessellation  $\mathcal{V}_\gamma^d \cap L_\ell$  a Voronoi tessellation?

**Chiu, Van De Weygaert, Stoyan, 1996:** "The sectional Poisson-Voronoi tessellation is not a Voronoi tessellation".

Given a  $\ell$ -dimensional affine subspace  $L_\ell \subset \mathbb{R}^d$  define the intersection tessellation

$$\mathcal{V}_\gamma^d \cap L_\ell := \{t \cap L_\ell : t \in \mathcal{V}_\gamma^d\}.$$

Useful formula:  $\mathbb{E}V_{d-\ell}(Z_d) = C_3(d, \ell) \gamma_\ell (\mathcal{V}_\gamma^d \cap L_\ell) \gamma^{-1}$ .

**Question:** Is the sectional Poisson-Voronoi tessellation  $\mathcal{V}_\gamma^d \cap L_\ell$  a Voronoi tessellation?

**Chiu, Van De Weygaert, Stoyan, 1996:** "The sectional Poisson-Voronoi tessellation is not a Voronoi tessellation".


**Next question:** How does the sectional Poisson-Voronoi tessellation look like?



# Poisson-Voronoi tessellation: graphical interpretation

Voronoi cell of (nuclei)  $v \in \eta$ :


$$V(v, \eta) := \{z \in \mathbb{R}^d : \|z - v\|^2 \leq \|z - v'\|^2 \text{ for all } v' \in \eta\}.$$



# Poisson-Voronoi tessellation: graphical interpretation

Voronoi cell of (nuclei)  $v \in \eta$ :


$$V(v, \eta) := \{z \in \mathbb{R}^d : \|z - v\|^2 \leq \|z - v'\|^2 \text{ for all } v' \in \eta\}.$$



# Poisson-Voronoi tessellation: graphical interpretation


Voronoi cell of (nuclei)  $v \in \eta$ :

$$V(v, \eta) := \{z \in \mathbb{R}^d : \|z - v\|^2 \leq \|z - v'\|^2 \text{ for all } v' \in \eta\}.$$




Voronoi cell of (nuclei)  $v \in \eta$ :

$$V(v, \eta) := \{z \in \mathbb{R}^d : \|z - v\|^2 \leq \|z - v'\|^2 \text{ for all } v' \in \eta\}.$$



Voronoi cell of (nuclei)  $v \in \eta$ :


$$V(v, \eta) := \{z \in \mathbb{R}^d : \|z - v\|^2 \leq \|z - v'\|^2 \text{ for all } v' \in \eta\}.$$



Voronoi cell of (nuclei)  $v \in \eta$ :

$$V(v, \eta) := \{z \in \mathbb{R}^d : \|z - v\|^2 \leq \|z - v'\|^2 \text{ for all } v' \in \eta\}.$$

- Cell  $V(v, \eta) \leftrightarrow$  paraboloid  $\Pi_{(v,0)}^d := \{(w, t) \in \mathbb{R}^{d+1} : t = \|v - w\|^2\}.$




# Poisson-Voronoi tessellation: graphical interpretation

Voronoi cell of (nuclei)  $v \in \eta$ :

$$V(v, \eta) := \{z \in \mathbb{R}^d : \|z - v\|^2 \leq \|z - v'\|^2 \text{ for all } v' \in \eta\}.$$


► Cell  $V(v, \eta) \leftrightarrow$  paraboloid  $\Pi_{(v,0)}^d$ .



Voronoi cell of (nuclei)  $v \in \eta$ :

$$V(v, \eta) := \{z \in \mathbb{R}^d : \|z - v\|^2 \leq \|z - v'\|^2 \text{ for all } v' \in \eta\}.$$

- Cell  $V(v, \eta) \leftrightarrow$  paraboloid  $\Pi_{(v,0)}^d$ .






Voronoi cell of (nuclei)  $v \in \eta$ :

$$V(v, \eta) := \{z \in \mathbb{R}^d : \|z - v\|^2 \leq \|z - v'\|^2 \text{ for all } v' \in \eta\}.$$


- ▶ Cell  $V(v, \eta) \leftrightarrow$  paraboloid  $\Pi_{(v,0)}^d$ .
- ▶  $\Pi_{(v,0)}^d \cap (L \times \mathbb{R}) \leftrightarrow \Pi_{(\bar{v},h)}^\ell$  in  $L \cong \mathbb{R}^\ell$ .



# Poisson-Voronoi tessellation: graphical interpretation

Generalized Voronoi cell of  $(v, h) \in \xi$ :


$$V((v, h), \xi) := \{z \in \mathbb{R}^d : \|z - v\|^2 + h \leq \|z - v'\|^2 + h' \text{ for all } (v', h') \in \xi\}.$$



# Poisson-Voronoi tessellation: graphical interpretation

Generalized Voronoi cell of  $(v, h) \in \xi$ :


$$V((v, h), \xi) := \{z \in \mathbb{R}^d : \|z - v\|^2 + h \leq \|z - v'\|^2 + h' \text{ for all } (v', h') \in \xi\}.$$



# Poisson-Voronoi tessellation: graphical interpretation

Generalized Voronoi cell of  $(v, h) \in \xi$ :


$$V((v, h), \xi) := \{z \in \mathbb{R}^d : \|z - v\|^2 + h \leq \|z - v'\|^2 + h' \text{ for all } (v', h') \in \xi\}.$$



# Generalized Voronoi tessellation (Laguerre tessellation)

Let  $\xi$  be a PPP in  $\mathbb{R}^d \times E$ ,  $E \subset \mathbb{R}$  with intensity measure having density of the form

$$(v, h) \mapsto \gamma f(h), \quad \gamma > 0.$$



# Generalized Voronoi tessellation (Laguerre tessellation)

Let  $\xi$  be a PPP in  $\mathbb{R}^d \times E$ ,  $E \subset \mathbb{R}$  with intensity measure having density of the form

$$(v, h) \mapsto \gamma f(h), \quad \gamma > 0.$$

**Laguerre cell** of  $(v, h) \in \xi$ :

$$V((v, h), \xi) := \{z \in \mathbb{R}^d : \|z - v\|^2 + h \leq \|z - v'\|^2 + h' \text{ for all } (v', h') \in \xi\}.$$


# Generalized Voronoi tessellation (Laguerre tessellation)

Let  $\xi$  be a PPP in  $\mathbb{R}^d \times E$ ,  $E \subset \mathbb{R}$  with intensity measure having density of the form

$$(v, h) \mapsto \gamma f(h), \quad \gamma > 0.$$

**Laguerre cell** of  $(v, h) \in \xi$ :

$$V((v, h), \xi) := \{z \in \mathbb{R}^d : \|z - v\|^2 + h \leq \|z - v'\|^2 + h' \text{ for all } (v', h') \in \xi\}.$$



# Generalized Voronoi tessellation (Laguerre tessellation)


Let  $\xi$  be a PPP in  $\mathbb{R}^d \times E$ ,  $E \subset \mathbb{R}$  with intensity measure having density of the form

$$(v, h) \mapsto \gamma f(h), \quad \gamma > 0.$$

**Laguerre cell** of  $(v, h) \in \xi$ :

$$V((v, h), \xi) := \{z \in \mathbb{R}^d : \|z - v\|^2 + h \leq \|z - v'\|^2 + h' \text{ for all } (v', h') \in \xi\}.$$

Denote  $\mathcal{L}(\xi) := \{V((v, h), \xi) : (v, h) \in \xi, \text{int} V((v, h), \xi) \neq \emptyset\}$ .





# Generalized Voronoi tessellation (Laguerre tessellation)

Let  $\xi$  be a PPP in  $\mathbb{R}^d \times E$ ,  $E \subset \mathbb{R}$  with intensity measure having density of the form

$$(v, h) \mapsto \gamma f(h), \quad \gamma > 0.$$

**Laguerre cell** of  $(v, h) \in \xi$ :

$$V((v, h), \xi) := \{z \in \mathbb{R}^d : \|z - v\|^2 + h \leq \|z - v'\|^2 + h' \text{ for all } (v', h') \in \xi\}.$$

Denote  $\mathcal{L}(\xi) := \{V((v, h), \xi) : (v, h) \in \xi, \text{int} V((v, h), \xi) \neq \emptyset\}$ .

# Generalized Voronoi tessellation (Laguerre tessellation)

Let  $\xi$  be a PPP in  $\mathbb{R}^d \times E$ ,  $E \subset \mathbb{R}$  with intensity measure having density of the form

$$(v, h) \mapsto \gamma f(h), \quad \gamma > 0.$$

**Laguerre cell** of  $(v, h) \in \xi$ :

$$V((v, h), \xi) := \{z \in \mathbb{R}^d : \|z - v\|^2 + h \leq \|z - v'\|^2 + h' \text{ for all } (v', h') \in \xi\}.$$


Denote  $\mathcal{L}(\xi) := \{V((v, h), \xi) : (v, h) \in \xi, \text{int} V((v, h), \xi) \neq \emptyset\}$ .

Our models:

►  $\xi = \eta_\beta$ ,  $\beta > -1$  is a PPP in  $\mathbb{R}^d \times [0, \infty)$  with

$$f(h) = c_{d+1, \beta} h^\beta, \quad c_{d+1, \beta} = \pi^{-\frac{d+1}{2}} \frac{\Gamma(\frac{d+3}{2} + \beta)}{\Gamma(\beta+1)}.$$

We denote  $\mathcal{V}_{\gamma, \beta}^d = \mathcal{L}(\eta_\beta)$ .



# Generalized Voronoi tessellation (Laguerre tessellation)

Let  $\xi$  be a PPP in  $\mathbb{R}^d \times E$ ,  $E \subset \mathbb{R}$  with intensity measure having density of the form

$$(v, h) \mapsto \gamma f(h), \quad \gamma > 0.$$

**Laguerre cell** of  $(v, h) \in \xi$ :

$$V((v, h), \xi) := \{z \in \mathbb{R}^d : \|z - v\|^2 + h \leq \|z - v'\|^2 + h' \text{ for all } (v', h') \in \xi\}.$$

Denote  $\mathcal{L}(\xi) := \{V((v, h), \xi) : (v, h) \in \xi, \text{int} V((v, h), \xi) \neq \emptyset\}$ .

Our models:

- $\xi = \eta_\beta$ ,  $\beta > -1$  is a PPP in  $\mathbb{R}^d \times [0, \infty)$  with


$$f(h) = c_{d+1, \beta} h^\beta.$$

We denote  $\mathcal{V}_{\gamma, \beta}^d = \mathcal{L}(\eta_\beta)$ .

- $\xi = \tilde{\eta}$  is a PPP in  $\mathbb{R}^d$  with

$$f(h) = e^{\lambda h}, \quad \lambda > 0.$$

We denote  $\tilde{\mathcal{V}}_\lambda^d = \mathcal{L}(\tilde{\eta})$ .



# Generalized Voronoi tessellation (Laguerre tessellation)

Let  $\xi$  be a PPP in  $\mathbb{R}^d \times E$ ,  $E \subset \mathbb{R}$  with intensity measure having density of the form

$$(v, h) \mapsto \gamma f(h), \quad \gamma > 0.$$

**Laguerre cell** of  $(v, h) \in \xi$ :

$$V((v, h), \xi) := \{z \in \mathbb{R}^d : \|z - v\|^2 + h \leq \|z - v'\|^2 + h' \text{ for all } (v', h') \in \xi\}.$$

Denote  $\mathcal{L}(\xi) := \{V((v, h), \xi) : (v, h) \in \xi, \text{int} V((v, h), \xi) \neq \emptyset\}$ .

Our models:

- $\xi = \eta_\beta$ ,  $\beta > -1$  is a PPP in  $\mathbb{R}^d \times [0, \infty)$  with


$$f(h) = c_{d+1, \beta} h^\beta.$$

We denote  $\mathcal{V}_{\gamma, \beta}^d = \mathcal{L}(\eta_\beta)$ .

- $\xi = \tilde{\eta}$  is a PPP in  $\mathbb{R}^d$  with

$$f(h) = e^{\lambda h}, \quad \lambda > 0.$$

We denote  $\tilde{\mathcal{V}}_\lambda^d = \mathcal{L}(\tilde{\eta})$ .



A.G., Z. Kabluchko and C. Thäle, 2022: The collections  $\mathcal{V}_{\gamma, \beta}^d$  and  $\tilde{\mathcal{V}}_\lambda^d$  are normal random tessellations and we call it  $\beta$ - and Gaussian-Voronoi tessellation, respectively.

## Theorem (A.G., Z. Kabluchko and C. Thäle, 2023)



For any  $\ell$ -dimensional affine subspace  $L_\ell$  we have

►  $\mathcal{V}_\gamma^d \cap L_\ell \stackrel{d}{=} \mathcal{V}_{\gamma_d, (d-\ell)/2-1}^\ell$  (up to isometry), where  $\gamma_d = \pi^{\frac{d+1}{2}} \Gamma(\frac{d+1}{2})^{-1} \gamma$ ;

## Theorem (A.G., Z. Kabluchko and C. Thäle, 2023)

For any  $\ell$ -dimensional affine subspace  $L_\ell$  we have



- $\mathcal{V}_\gamma^d \cap L_\ell \stackrel{d}{=} \mathcal{V}_{\gamma_d, (d-\ell)/2-1}^\ell$  (up to isometry), where  $\gamma_d = \pi^{\frac{d+1}{2}} \Gamma(\frac{d+1}{2})^{-1} \gamma$ ;




## Theorem (A.G., Z. Kabluchko and C. Thäle, 2023)

For any  $\ell$ -dimensional affine subspace  $L_\ell$  we have

- ▶  $\mathcal{V}_\gamma^d \cap L_\ell \stackrel{d}{=} \mathcal{V}_{\gamma_d, (d-\ell)/2-1}^\ell$  (up to isometry), where  $\gamma_d = \pi^{\frac{d+1}{2}} \Gamma(\frac{d+1}{2})^{-1} \gamma$ ;
- ▶  $\mathcal{V}_{\gamma, \beta}^d \cap L_\ell \stackrel{d}{=} \mathcal{V}_{\gamma, (d-\ell)/2+\beta}^\ell$  (up to isometry) for any  $\beta > -1$ .




## $\beta$ -Delaunay tessellation (definition via duality)



$\beta$ -Delaunay tessellation  $\mathcal{D}_{\gamma,\beta}^d$  is **dual** of  $\beta$ -Voronoi tessellation of  $\mathcal{V}_{\gamma,\beta}^d$ .




## $\beta$ -Delaunay tessellation (definition via duality)



$\beta$ -Delaunay tessellation  $\mathcal{D}_{\gamma,\beta}^d$  is **dual** of  $\beta$ -Voronoi tessellation of  $\mathcal{V}_{\gamma,\beta}^d$ .

► Center of cell  $z(V((v, h), \eta_\beta)) = v$ .


## $\beta$ -Delaunay tessellation (definition via duality)



$\beta$ -Delaunay tessellation  $\mathcal{D}_{\gamma,\beta}^d$  is **dual** of  $\beta$ -Voronoi tessellation of  $\mathcal{V}_{\gamma,\beta}^d$ .

- ▶ Center of cell  $z(V((v, h), \eta_\beta)) = v$ .
- ▶ Define  $D(y) := \text{conv}\{v : y\text{-vertex of } V((v, h), \eta_\beta)\}$ .


## $\beta$ -Delaunay tessellation (definition via duality)



$\beta$ -Delaunay tessellation  $\mathcal{D}_{\gamma,\beta}^d$  is **dual** of  $\beta$ -Voronoi tessellation of  $\mathcal{V}_{\gamma,\beta}^d$ .

- ▶ Center of cell  $z(V((v, h), \eta_\beta)) = v$ .
- ▶ Define  $D(y) := \text{conv}\{v : y\text{-vertex of } V((v, h), \eta_\beta)\}$ .


## $\beta$ -Delaunay tessellation (definition via duality)



$\beta$ -Delaunay tessellation  $\mathcal{D}_{\gamma,\beta}^d$  is **dual** of  $\beta$ -Voronoi tessellation of  $\mathcal{V}_{\gamma,\beta}^d$ .

- ▶ Center of cell  $z(V((v, h), \eta_\beta)) = v$ .
- ▶ Define  $D(y) := \text{conv}\{v : y\text{-vertex of } V((v, h), \eta_\beta)\}$ .
- ▶  $\mathcal{D}_{\gamma,\beta}^d = \{D(y), y \in \mathcal{F}_0(\mathcal{V}_{\gamma,\beta}^d)\}$ .

## $\beta$ -Delaunay tessellation (definition via duality)



$\beta$ -Delaunay tessellation  $\mathcal{D}_{\gamma,\beta}^d$  is **dual** of  $\beta$ -Voronoi tessellation of  $\mathcal{V}_{\gamma,\beta}^d$ .

**Fact:**  $\mathcal{D}_{\beta}$  is almost surely a simplicial random tessellations.

Theorem (A.G., Z. Kabluchko and C.Thäle, 2022)

We have  $Z(\mathcal{D}_{\gamma,\beta}^d) \stackrel{d}{=} R \cdot \text{conv}(Y_1, \dots, Y_{d+1})$ , where

(a)  $(Y_1, \dots, Y_{d+1})$  are random points, whose joint density is

$$(y_1, \dots, y_{d+1}) \mapsto \text{const} \cdot \text{Vol}(\text{conv}(y_1, \dots, y_{d+1})) \prod_{i=1}^{d+1} (1 - \|y_i\|^2)^\beta \mathbf{1}_{\mathbb{B}^d}(\|y_i\|);$$

(b)  $R$  is a random variable on  $(0, \infty)$  independent of  $(Y_1, \dots, Y_{d+1})$  with density:

$$r \mapsto \text{const} \cdot r^{(d+1)(d+1+2\beta)} e^{-\gamma \cdot \text{const} \cdot r^{d+2+2\beta}}.$$

## Theorem (A.G., Z. Kabluchko and C.Thäle, 2022)

We have  $Z(\mathcal{D}_{\gamma,\beta}^d) \stackrel{d}{=} R \cdot \text{conv}(Y_1, \dots, Y_{d+1})$ , where

(a)  $(Y_1, \dots, Y_{d+1})$  are random points, whose joint density is

$$(y_1, \dots, y_{d+1}) \mapsto \text{const} \cdot \text{Vol}(\text{conv}(y_1, \dots, y_{d+1})) \prod_{i=1}^{d+1} (1 - \|y_i\|^2)^\beta \mathbf{1}_{\mathbb{B}^d}(\|y_i\|);$$

(b)  $R$  is a random variable on  $(0, \infty)$  independent of  $(Y_1, \dots, Y_{d+1})$  with density:

$$r \mapsto \text{const} \cdot r^{(d+1)(d+1+2\beta)} e^{-\gamma \cdot \text{const} \cdot r^{d+2+2\beta}}.$$

Remark:  $Z(\mathcal{D}_{\gamma,\beta}^d)$  is randomly rescaled volume-weighted beta simplices;

► Recall:  $\mathcal{D}_{\gamma,\beta}^d = \{D(y) : y \in \mathcal{F}_0(\mathcal{V}_{\gamma,\beta}^d)\}$

$$D(y) = \text{conv}\{v_1, \dots, v_{d+1} : y\text{-vertex of } V((v_i, h_i), \eta_\beta), 1 \leq i \leq d+1\}$$




# Alternative definition of $\mathcal{D}_{\gamma,\beta}^d$

► Recall:  $\mathcal{D}_{\gamma,\beta}^d = \{D(y) : y \in \mathcal{F}_0(\mathcal{V}_{\gamma,\beta}^d)\}$

$$D(y) = \text{conv}\{v_1, \dots, v_{d+1} : y\text{-vertex of } V((v_i, h_i), \eta_\beta), 1 \leq i \leq d+1\}$$

$\iff$  there is a constant  $h_y$  s.t. a standard downward paraboloid with apex  $(y, h_y)$  contains points  $(v_i, h_i)$ ,  $1 \leq i \leq d+1$  on its boundary and no other points of  $\eta_\beta$  below its surface.




# Alternative definition of $\mathcal{D}_{\gamma,\beta}^d$

► Recall:  $\mathcal{D}_{\gamma,\beta}^d = \{D(y) : y \in \mathcal{F}_0(\mathcal{V}_{\gamma,\beta}^d)\}$

$$D(y) = \text{conv}\{v_1, \dots, v_{d+1} : y\text{- vertex of } V((v_i, h_i), \eta_\beta), 1 \leq i \leq d+1\}$$

$\iff$  there is a constant  $h_y$  s.t. a standard downward paraboloid with apex  $(y, h_y)$  contains points  $(v_i, h_i)$ ,  $1 \leq i \leq d+1$  on its boundary and no other points of  $\eta_\beta$  below its surface.




# Alternative definition of $\mathcal{D}_{\gamma,\beta}^d$

► Recall:  $\mathcal{D}_{\gamma,\beta}^d = \{D(y) : y \in \mathcal{F}_0(\mathcal{V}_{\gamma,\beta}^d)\}$

$$D(y) = \text{conv}\{v_1, \dots, v_{d+1} : y\text{-vertex of } V((v_i, h_i), \eta_\beta), 1 \leq i \leq d+1\}$$

$\iff$  there is a constant  $h_y$  s.t. a standard downward paraboloid with apex  $(y, h_y)$  contains points  $(v_i, h_i)$ ,  $1 \leq i \leq d+1$  on its boundary and no other points of  $\eta_\beta$  below its surface.




## Alternative definition of $\mathcal{D}_{\gamma,\beta}^d$

► Recall:  $\mathcal{D}_{\gamma,\beta}^d = \{D(y) : y \in \mathcal{F}_0(\mathcal{V}_{\gamma,\beta}^d)\}$

$$D(y) = \text{conv}\{v_1, \dots, v_{d+1} : y\text{-vertex of } V((v_i, h_i), \eta_\beta), 1 \leq i \leq d+1\}$$

$\iff$  there is a constant  $h_y$  s.t. a standard downward paraboloid with apex  $(y, h_y)$  contains points  $(v_i, h_i)$ ,  $1 \leq i \leq d+1$  on its boundary and no other points of  $\eta_\beta$  below its surface.




## Alternative definition of $\mathcal{D}_{\gamma,\beta}^d$

► Recall:  $\mathcal{D}_{\gamma,\beta}^d = \{D(y) : y \in \mathcal{F}_0(\mathcal{V}_{\gamma,\beta}^d)\}$

$$D(y) = \text{conv}\{v_1, \dots, v_{d+1} : y\text{-vertex of } V((v_i, h_i), \eta_\beta), 1 \leq i \leq d+1\}$$

$\iff$  there is a constant  $h_y$  s.t. a standard downward paraboloid with apex  $(y, h_y)$  contains points  $(v_i, h_i)$ ,  $1 \leq i \leq d+1$  on its boundary and no other points of  $\eta_\beta$  below its surface.



# Expected intrinsic volumes of the typical cell of Poisson-Voronoi tessellation

As a consequence of exact stochastic representation we have:

As a consequence of exact stochastic representation we have:

- Explicit formulas for  $\mathbb{E}V_d(Z(\mathcal{D}_{\gamma,\beta}^d))$  (via Gamma functions);

# Expected intrinsic volumes of the typical cell of Poisson-Voronoi tessellation

As a consequence of exact stochastic representation we have:

- ▶ Explicit formulas for  $\mathbb{E}V_d(Z(\mathcal{D}_{\gamma,\beta}^d))$  (via Gamma functions);
- ▶ Let

$$\sigma_0(P) = \sum_{y \in \mathcal{F}_0(P)} \alpha(y, P),$$

where  $\alpha(y, P)$  is the solid angle of  $P$  at  $y$ .



# Expected intrinsic volumes of the typical cell of Poisson-Voronoi tessellation


As a consequence of exact stochastic representation we have:

- Explicit formulas for  $\mathbb{E}V_d(Z(\mathcal{D}_{\gamma,\beta}^d))$  (via Gamma functions);
- Let

$$\sigma_0(P) = \sum_{y \in \mathcal{F}_0(P)} \alpha(y, P),$$

where  $\alpha(y, P)$  is the solid angle of  $P$  at  $y$ . Then

$$\mathbb{E}\sigma_0(Z(\mathcal{D}_{\gamma,\beta}^d)) = \binom{d+1}{k+1} J_{d+1,1}(\beta + \frac{1}{2}).$$



# Expected intrinsic volumes of the typical cell of Poisson-Voronoi tessellation

As a consequence of exact stochastic representation we have:


- Explicit formulas for  $\mathbb{E}V_d(Z(\mathcal{D}_{\gamma,\beta}^d))$  (via Gamma functions);
- Let

$$\sigma_0(P) = \sum_{y \in \mathcal{F}_0(P)} \alpha(y, P),$$

where  $\alpha(y, P)$  is the solid angle of  $P$  at  $y$ . Then

$$\mathbb{E}\sigma_0(Z(\mathcal{D}_{\gamma,\beta}^d)) = \binom{d+1}{k+1} J_{d+1,1}(\beta + \frac{1}{2}).$$

- $\gamma_0(\mathcal{D}_{\gamma,\beta}^d) = \gamma_d(\mathcal{D}_{\gamma,\beta}^d) \mathbb{E}\sigma_0(Z(\mathcal{D}_{\gamma,\beta}^d))$ .



# Expected intrinsic volumes of the typical cell of Poisson-Voronoi tessellation

As a consequence of exact stochastic representation we have:


- Explicit formulas for  $\mathbb{E}V_d(Z(\mathcal{D}_{\gamma,\beta}^d))$  (via Gamma functions);
- Let

$$\sigma_0(P) = \sum_{y \in \mathcal{F}_0(P)} \alpha(y, P),$$

where  $\alpha(y, P)$  is the solid angle of  $P$  at  $y$ . Then

$$\mathbb{E}\sigma_0(Z(\mathcal{D}_{\gamma,\beta}^d)) = \binom{d+1}{k+1} J_{d+1,1}(\beta + \frac{1}{2}).$$

- $\gamma_0(\mathcal{D}_{\gamma,\beta}^d) = [\mathbb{E}V_d(Z(\mathcal{D}_{\gamma,\beta}^d))]^{-1} \mathbb{E}\sigma_0(Z(\mathcal{D}_{\gamma,\beta}^d)).$



# Expected intrinsic volumes of the typical cell of Poisson-Voronoi tessellation

As a consequence of exact stochastic representation we have:


- Explicit formulas for  $\mathbb{E}V_d(Z(\mathcal{D}_{\gamma,\beta}^d))$  (via Gamma functions);
- Let

$$\sigma_0(P) = \sum_{y \in \mathcal{F}_0(P)} \alpha(y, P),$$

where  $\alpha(y, P)$  is the solid angle of  $P$  at  $y$ . Then

$$\mathbb{E}\sigma_0(Z(\mathcal{D}_{\gamma,\beta}^d)) = \binom{d+1}{k+1} J_{d+1,1}(\beta + \tfrac{1}{2}).$$

- $\gamma_0(\mathcal{D}_{\gamma,\beta}^d) = [\mathbb{E}V_d(Z(\mathcal{D}_{\gamma,\beta}^d))]^{-1} \mathbb{E}\sigma_0(Z(\mathcal{D}_{\gamma,\beta}^d)) = \gamma_d(\mathcal{V}_{\gamma,\beta}^d).$



# Expected intrinsic volumes of the typical cell of Poisson-Voronoi tessellation

As a consequence of exact stochastic representation we have:


- Explicit formulas for  $\mathbb{E}V_d(Z(\mathcal{D}_{\gamma,\beta}^d))$  (via Gamma functions);
- Let

$$\sigma_0(P) = \sum_{y \in \mathcal{F}_0(P)} \alpha(y, P),$$

where  $\alpha(y, P)$  is the solid angle of  $P$  at  $y$ . Then

$$\mathbb{E}\sigma_0(Z(\mathcal{D}_{\gamma,\beta}^d)) = \binom{d+1}{k+1} J_{d+1,1}(\beta + \frac{1}{2}).$$

- $\gamma_0(\mathcal{D}_{\gamma,\beta}^d) = [\mathbb{E}V_d(Z(\mathcal{D}_{\gamma,\beta}^d))]^{-1} \mathbb{E}\sigma_0(Z(\mathcal{D}_{\gamma,\beta}^d)) = \gamma_d(\mathcal{V}_{\gamma,\beta}^d).$



Corollary (A.G., Z. Kabluchko and C. Thäle, 2023)

For any  $0 \leq j \leq d$  we have

$$\mathbb{E}V_j(Z(\mathcal{V}_{\gamma}^d)) = \gamma^{-\frac{j}{d}} J_{d-j+1,1}\left(\frac{j-1}{2}\right) \frac{2^{d-j+1} \pi^{\frac{d-j}{2}}}{d(d-j)!} \frac{\Gamma(\frac{(d-j+1)(d-1)}{2} + 1)}{\Gamma(\frac{(d-j+1)(d-1)+1}{2})} \frac{\Gamma(\frac{d}{2} + 1)^{d-j+\frac{j}{d}}}{\Gamma(\frac{d+1}{2})^{d-j}} \frac{\Gamma(d-j+\frac{j}{d})}{\Gamma(\frac{j+1}{2})}.$$

# Expected intrinsic volumes of the typical cell of Poisson-Voronoi tessellation

As a consequence of exact stochastic representation we have:


- Explicit formulas for  $\mathbb{E}V_d(Z(\mathcal{D}_{\gamma,\beta}^d))$  (via Gamma functions);
- Let

$$\sigma_0(P) = \sum_{y \in \mathcal{F}_0(P)} \alpha(y, P),$$

where  $\alpha(y, P)$  is the solid angle of  $P$  at  $y$ . Then

$$\mathbb{E}\sigma_0(Z(\mathcal{D}_{\gamma,\beta}^d)) = \binom{d+1}{k+1} J_{d+1,1} \left( \beta + \frac{1}{2} \right).$$

- $\gamma_0(\mathcal{D}_{\gamma,\beta}^d) = [\mathbb{E}V_d(Z(\mathcal{D}_{\gamma,\beta}^d))]^{-1} \mathbb{E}\sigma_0(Z(\mathcal{D}_{\gamma,\beta}^d)) = \gamma_d(\mathcal{V}_{\gamma,\beta}^d).$



## Corollary (A.G., Z. Kabluchko and C. Thäle, 2023)

For any  $0 \leq j \leq d$  we have

$$\mathbb{E}V_j(Z(\mathcal{V}_{\gamma}^d)) = \gamma^{-\frac{j}{d}} J_{d-j+1,1} \left( \frac{j-1}{2} \right) \frac{2^{d-j+1} \pi^{\frac{d-j}{2}}}{d(d-j)!} \frac{\Gamma(\frac{(d-j+1)(d-1)}{2} + 1)}{\Gamma(\frac{(d-j+1)(d-1)+1}{2})} \frac{\Gamma(\frac{d}{2} + 1)^{d-j+\frac{j}{d}}}{\Gamma(\frac{d+1}{2})^{d-j}} \frac{\Gamma(d-j+\frac{j}{d})}{\Gamma(\frac{j+1}{2})}.$$

$$\mathbb{E}V_j(Z(\mathcal{V}_{\gamma}^d)) \Leftarrow \gamma_{d-j}(\mathcal{V}_{\gamma}^d \cap L_{d-j}) \Leftarrow \gamma_{d-j}(\mathcal{V}_{\gamma_{d,j/2-1}}^{d-j})$$

## Corollary

For any  $d \geq 2$ ,  $1 \leq \ell \leq d - 1$  and  $L_\ell \in A(d, \ell)$  we have

$$\mathbb{E} V_\ell(Z(\mathcal{V}_\gamma^d \cap L_\ell)) = \frac{d\gamma^{-\frac{\ell}{d}}}{2J_{\ell+1,1}(\frac{d-\ell-1}{2})\pi^{\frac{\ell}{2}}} \frac{\Gamma(\frac{(\ell+1)(d-1)+1}{2})}{\Gamma(\frac{(\ell+1)(d-1)}{2} + 1)} \frac{\Gamma(\frac{\ell+2}{2})}{\Gamma(\ell + 1 - \frac{\ell}{d})} \frac{\Gamma(\frac{d+1}{2})^{\ell+1}}{\Gamma(\frac{d}{2} + 1)^{\ell+1-\frac{\ell}{d}}}.$$

## Corollary

For any  $d \geq 2$ ,  $1 \leq \ell \leq d - 1$  and  $L_\ell \in A(d, \ell)$  we have

$$\mathbb{E} V_\ell(Z(\mathcal{V}_\gamma^d \cap L_\ell)) = \frac{d\gamma^{-\frac{\ell}{d}}}{2J_{\ell+1,1}(\frac{d-\ell-1}{2})\pi^{\frac{\ell}{2}}} \frac{\Gamma(\frac{(\ell+1)(d-1)+1}{2})}{\Gamma(\frac{(\ell+1)(d-1)}{2} + 1)} \frac{\Gamma(\frac{\ell+2}{2})}{\Gamma(\ell + 1 - \frac{\ell}{d})} \frac{\Gamma(\frac{d+1}{2})^{\ell+1}}{\Gamma(\frac{d}{2} + 1)^{\ell+1-\frac{\ell}{d}}}.$$

## Corollary

For any  $\ell \in \mathbb{N}$  and  $L_\ell \in A(d, \ell)$  we have

$$\lim_{d \rightarrow \infty} \mathbb{E} V_\ell(Z(\mathcal{V}_\gamma^d \cap L_\ell)) = (\pi e)^{-\frac{\ell}{2}} J_{\ell+1,1}(\infty)^{-1} \frac{\Gamma(\frac{\ell}{2})}{2(\ell-1)!\sqrt{\ell+1}},$$

where  $J_{\ell+1,1}(\infty)$  is a solid angle of a regular  $\ell$ -dimensional simplex.



## Corollary

For any  $d \geq 2$ ,  $1 \leq \ell \leq d - 1$  and  $L_\ell \in A(d, \ell)$  we have

$$\mathbb{E}V_\ell(Z(\mathcal{V}_\gamma^d \cap L_\ell)) = \frac{d\gamma^{-\frac{\ell}{d}}}{2J_{\ell+1,1}(\frac{d-\ell-1}{2})\pi^{\frac{\ell}{2}}} \frac{\Gamma(\frac{(\ell+1)(d-1)+1}{2})}{\Gamma(\frac{(\ell+1)(d-1)}{2} + 1)} \frac{\Gamma(\frac{\ell+2}{2})}{\Gamma(\ell + 1 - \frac{\ell}{d})} \frac{\Gamma(\frac{d+1}{2})^{\ell+1}}{\Gamma(\frac{d}{2} + 1)^{\ell+1-\frac{\ell}{d}}}.$$

## Corollary

For any  $\ell \in \mathbb{N}$  and  $L_\ell \in A(d, \ell)$  we have

$$\lim_{d \rightarrow \infty} \mathbb{E}V_\ell(Z(\mathcal{V}_\gamma^d \cap L_\ell)) = (\pi e)^{-\frac{\ell}{2}} J_{\ell+1,1}(\infty)^{-1} \frac{\Gamma(\frac{\ell}{2})}{2(\ell-1)!\sqrt{\ell+1}},$$

where  $J_{\ell+1,1}(\infty)$  is a solid angle of a regular  $\ell$ -dimensional simplex.

**Remark:** Case  $\ell = 1, 2$  was obtained by [Miles, 1984](#).

## Corollary

For any  $d \geq 2$ ,  $1 \leq \ell \leq d - 1$  and  $L_\ell \in A(d, \ell)$  we have

$$\mathbb{E}V_\ell(Z(\mathcal{V}_\gamma^d \cap L_\ell)) = \frac{d\gamma^{-\frac{\ell}{d}}}{2J_{\ell+1,1}(\frac{d-\ell-1}{2})\pi^{\frac{\ell}{2}}} \frac{\Gamma(\frac{(\ell+1)(d-1)+1}{2})}{\Gamma(\frac{(\ell+1)(d-1)}{2} + 1)} \frac{\Gamma(\frac{\ell+2}{2})}{\Gamma(\ell + 1 - \frac{\ell}{d})} \frac{\Gamma(\frac{d+1}{2})^{\ell+1}}{\Gamma(\frac{d}{2} + 1)^{\ell+1-\frac{\ell}{d}}}.$$

## Corollary

For any  $\ell \in \mathbb{N}$  and  $L_\ell \in A(d, \ell)$  we have

$$\lim_{d \rightarrow \infty} \mathbb{E}V_1(Z(\mathcal{V}_\gamma^d \cap L_\ell)) = (2e)^{-\frac{1}{2}},$$

where  $J_{\ell+1,1}(\infty)$  is a solid angle of a regular  $\ell$ -dimensional simplex.

**Remark:** Case  $\ell = 1, 2$  was obtained by [Miles, 1984](#).

## Corollary

For any  $d \geq 2$ ,  $1 \leq \ell \leq d - 1$  and  $L_\ell \in A(d, \ell)$  we have

$$\mathbb{E} V_\ell(Z(\mathcal{V}_\gamma^d \cap L_\ell)) = \frac{d\gamma^{-\frac{\ell}{d}}}{2J_{\ell+1,1}(\frac{d-\ell-1}{2})\pi^{\frac{\ell}{2}}} \frac{\Gamma(\frac{(\ell+1)(d-1)+1}{2})}{\Gamma(\frac{(\ell+1)(d-1)}{2} + 1)} \frac{\Gamma(\frac{\ell+2}{2})}{\Gamma(\ell + 1 - \frac{\ell}{d})} \frac{\Gamma(\frac{d+1}{2})^{\ell+1}}{\Gamma(\frac{d}{2} + 1)^{\ell+1-\frac{\ell}{d}}}.$$

## Corollary

For any  $\ell \in \mathbb{N}$  and  $L_\ell \in A(d, \ell)$  we have

$$\lim_{d \rightarrow \infty} \mathbb{E} V_2(Z(\mathcal{V}_\gamma^d \cap L_\ell)) = \sqrt{3}(\pi e)^{-1},$$

where  $J_{\ell+1,1}(\infty)$  is a solid angle of a regular  $\ell$ -dimensional simplex.

**Remark:** Case  $\ell = 1, 2$  was obtained by [Miles, 1984](#).

## Corollary

For any  $d \geq 2$ ,  $1 \leq \ell \leq d - 1$  and  $L_\ell \in A(d, \ell)$  we have

$$\mathbb{E} V_\ell(Z(\mathcal{V}_\gamma^d \cap L_\ell)) = \frac{d\gamma^{-\frac{\ell}{d}}}{2J_{\ell+1,1}(\frac{d-\ell-1}{2})\pi^{\frac{\ell}{2}}} \frac{\Gamma(\frac{(\ell+1)(d-1)+1}{2})}{\Gamma(\frac{(\ell+1)(d-1)}{2} + 1)} \frac{\Gamma(\frac{\ell+2}{2})}{\Gamma(\ell + 1 - \frac{\ell}{d})} \frac{\Gamma(\frac{d+1}{2})^{\ell+1}}{\Gamma(\frac{d}{2} + 1)^{\ell+1-\frac{\ell}{d}}}.$$

## Corollary

For any  $\ell \in \mathbb{N}$  and  $L_\ell \in A(d, \ell)$  we have

$$\lim_{d \rightarrow \infty} \mathbb{E} V_\ell(Z(\mathcal{V}_\gamma^d \cap L_\ell)) = (\pi e)^{-\frac{\ell}{2}} J_{\ell+1,1}(\infty)^{-1} \frac{\Gamma(\frac{\ell}{2})}{2(\ell-1)!\sqrt{\ell+1}},$$

where  $J_{\ell+1,1}(\infty)$  is a solid angle of a regular  $\ell$ -dimensional simplex.

**Remark:** Case  $\ell = 1, 2$  was obtained by [Miles, 1984](#).


**Observation:**  $\lim_{d \rightarrow \infty} \mathbb{E} V_\ell(Z(\mathcal{V}_\gamma^d \cap L_\ell)) = \mathbb{E} V_\ell(Z(\tilde{\mathcal{V}}_{\pi e}^\ell)).$

## Theorem (A.G., Z. Kabluchko and C.Thäle, 2023)

For any  $\ell \in \mathbb{N}$ ,  $L_\ell \in A(d, \ell)$  and every ball  $B_R \subset \mathbb{R}^d$  of radius  $R > 0$  centered at the origin we have

$$\lim_{d \rightarrow \infty} \mathbb{P} \left[ \bigcup_{t \in (\mathcal{V}_\gamma^d \cap L_\ell)} (\text{bd } t \cap B_R) = \bigcup_{t \in \tilde{\mathcal{V}}_{\pi e}^\ell} (\text{bd } t \cap B_R) \right] = 1.$$

And  $Z(\mathcal{V}_\gamma^d \cap L_\ell) \xrightarrow{d} Z(\tilde{\mathcal{V}}_{\pi e}^\ell)$  as  $d \rightarrow \infty$ .



## Theorem (A.G., Z. Kabluchko and C.Thäle, 2023)

For any  $\ell \in \mathbb{N}$ ,  $L_\ell \in A(d, \ell)$  and every ball  $B_R \subset \mathbb{R}^d$  of radius  $R > 0$  centered at the origin we have

$$\lim_{d \rightarrow \infty} \mathbb{P} \left[ \bigcup_{t \in (\mathcal{V}_\gamma^d \cap L_\ell)} (\text{bd } t \cap B_R) = \bigcup_{t \in \tilde{\mathcal{V}}_{\pi e}^\ell} (\text{bd } t \cap B_R) \right] = 1.$$

And  $Z(\mathcal{V}_\gamma^d \cap L_\ell) \xrightarrow{d} Z(\tilde{\mathcal{V}}_{\pi e}^\ell)$  as  $d \rightarrow \infty$ .



- ▶ A. Gusakova, Z. Kabluchko and C. Thäle. **The  $\beta$ -Delaunay tessellation I: Description of the model and geometry of typical cells** - Journal of Applied Probability, 2022.
- ▶ A. Gusakova, Z. Kabluchko and C. Thäle. **The  $\beta$ -Delaunay tessellation II: The Gaussian limit tessellation** - Electronic Journal of Probability, 2022.
- ▶ A. Gusakova, Z. Kabluchko and C. Thäle. **Sectional Voronoi tessellations: Characterization and high-dimensional limits** - To appear in Bernoulli, 2023.

Thank you for attention!