

# Brunn-Minkowski inequalities for variational functionals in Gauss space

Andrea Colesanti

**“Convex geometry and geometric probability”**

Paris Lodron Universität - Salzburg

September 25-29, 2023

It was twenty years ago today...



Cortona, summer 2003

The original results presented in this talk were obtained in collaboration with: **Elisa Francini**, **Galyna Livshyts** and **Paolo Salani**.

# The Brunn-Minkowski inequality

# The Brunn-Minkowski inequality

**Thm.** *Let  $K_0, K_1$  be convex bodies in  $\mathbb{R}^n$  and let  $t \in [0, 1]$ ; then*

$$V_n((1-t)K_0 + tK_1)^{1/n} \geq (1-t)V_n(K_0)^{1/n} + tV_n(K_1)^{1/n}. \quad (\text{BM})$$

# The Brunn-Minkowski inequality

**Thm.** *Let  $K_0, K_1$  be convex bodies in  $\mathbb{R}^n$  and let  $t \in [0, 1]$ ; then*

$$V_n((1-t)K_0 + tK_1)^{1/n} \geq (1-t)V_n(K_0)^{1/n} + tV_n(K_1)^{1/n}. \quad (\text{BM})$$

*Equality holds if and only if  $K_0$  and  $K_1$  are homothetic, or they lie in parallel hyperplanes.*

# The Brunn-Minkowski inequality

**Thm.** Let  $K_0, K_1$  be convex bodies in  $\mathbb{R}^n$  and let  $t \in [0, 1]$ ; then

$$V_n((1-t)K_0 + tK_1)^{1/n} \geq (1-t)V_n(K_0)^{1/n} + tV_n(K_1)^{1/n}. \quad (\text{BM})$$

*Equality holds if and only if  $K_0$  and  $K_1$  are homothetic, or they lie in parallel hyperplanes.*

Here:

▶  $V_n = \text{volume (Lebesgue measure)}$ ;

▶

$$(1-t)K_0 + tK_1 = \{(1-t)x + ty : x \in K_0, y \in K_1\}.$$

# The Brunn-Minkowski inequality

**Thm.** Let  $K_0, K_1$  be convex bodies in  $\mathbb{R}^n$  and let  $t \in [0, 1]$ ; then

$$V_n((1-t)K_0 + tK_1)^{1/n} \geq (1-t)V_n(K_0)^{1/n} + tV_n(K_1)^{1/n}. \quad (\text{BM})$$

*Equality holds if and only if  $K_0$  and  $K_1$  are homothetic, or they lie in parallel hyperplanes.*

Here:

▶  $V_n = \text{volume (Lebesgue measure)}$ ;



$$(1-t)K_0 + tK_1 = \{(1-t)x + ty : x \in K_0, y \in K_1\}.$$

Equivalently: the functional  $V_n^{1/n}$  is concave in the class of convex bodies of  $\mathbb{R}^n$ , equipped with the Minkowski addition.



# Links with other inequalities

## Links with other inequalities

- ▶ The Brunn-Minkowski inequality provides a simple proof of the **isoperimetric inequality**.

## Links with other inequalities

- ▶ The Brunn-Minkowski inequality provides a simple proof of the **isoperimetric inequality**.
- ▶ The Brunn-Minkowski inequality is naturally connected to the **Prékopa-Leindler inequality**,

## Links with other inequalities

- ▶ The Brunn-Minkowski inequality provides a simple proof of the **isoperimetric inequality**.
- ▶ The Brunn-Minkowski inequality is naturally connected to the **Prékopa-Leindler inequality**, which represents its functional counterpart and gives an easy proof of it.

## Links with other inequalities

- ▶ The Brunn-Minkowski inequality provides a simple proof of the **isoperimetric inequality**.
- ▶ The Brunn-Minkowski inequality is naturally connected to the **Prékopa-Leindler inequality**, which represents its functional counterpart and gives an easy proof of it.
- ▶ The Brunn-Minkowski inequality is a special case of the family of **Alexandrov-Fenchel** inequalities.

# Links with other inequalities

- ▶ The Brunn-Minkowski inequality provides a simple proof of the **isoperimetric inequality**.
- ▶ The Brunn-Minkowski inequality is naturally connected to the **Prékopa-Leindler inequality**, which represents its functional counterpart and gives an easy proof of it.
- ▶ The Brunn-Minkowski inequality is a special case of the family of **Alexandrov-Fenchel** inequalities.

⋮

# Links with other inequalities

- ▶ The Brunn-Minkowski inequality provides a simple proof of the **isoperimetric inequality**.
- ▶ The Brunn-Minkowski inequality is naturally connected to the **Prékopa-Leindler inequality**, which represents its functional counterpart and gives an easy proof of it.
- ▶ The Brunn-Minkowski inequality is a special case of the family of **Alexandrov-Fenchel** inequalities.
- ▶  $\vdots$
- ▶ Much more about this topic can be found in: R. J. Gardner, *The Brunn-Minkowski inequality*, 2002.

# A Brunn-Minkowski type inequality for intrinsic volumes



# A Brunn-Minkowski type inequality for intrinsic volumes

Given a convex body  $K$ , let

$$V_0(K), V_1(K), \dots, V_n(K)$$

denote its *intrinsic volumes*.

## A Brunn-Minkowski type inequality for intrinsic volumes

Given a convex body  $K$ , let

$$V_0(K), V_1(K), \dots, V_n(K)$$

denote its *intrinsic volumes*.

As a consequence of the Alexandrov-Fenchel inequalities, we obtain, for  $i \geq 1$ :

$$V_i((1-t)K_0 + tK_1)^{1/i} \geq (1-t)V_i(K_0)^{1/i} + tV_i(K_1)^{1/i}$$

for every  $K_0, K_1 \in \mathcal{K}^n$ , for every  $t \in [0, 1]$ .

# A Brunn-Minkowski type inequality for intrinsic volumes

Given a convex body  $K$ , let

$$V_0(K), V_1(K), \dots, V_n(K)$$

denote its *intrinsic volumes*.

As a consequence of the Alexandrov-Fenchel inequalities, we obtain, for  $i \geq 1$ :

$$V_i((1-t)K_0 + tK_1)^{1/i} \geq (1-t)V_i(K_0)^{1/i} + tV_i(K_1)^{1/i}$$

for every  $K_0, K_1 \in \mathcal{K}^n$ , for every  $t \in [0, 1]$ .

Note that  $V_i$  is  $i$ -homogeneous:

$$V_i(sK) = s^i V_i(K) \quad \forall K \in \mathcal{K}^n, s > 0.$$

# A Brunn-Minkowski type inequality for intrinsic volumes

Given a convex body  $K$ , let

$$V_0(K), V_1(K), \dots, V_n(K)$$

denote its *intrinsic volumes*.

As a consequence of the Alexandrov-Fenchel inequalities, we obtain, for  $i \geq 1$ :

$$V_i((1-t)K_0 + tK_1)^{1/i} \geq (1-t)V_i(K_0)^{1/i} + tV_i(K_1)^{1/i}$$

for every  $K_0, K_1 \in \mathcal{K}^n$ , for every  $t \in [0, 1]$ .

Note that  $V_i$  is  $i$ -homogeneous:

$$V_i(sK) = s^i V_i(K) \quad \forall K \in \mathcal{K}^n, s > 0.$$

Hence, as for the volume,  $V_i$  raised to the reciprocal of its homogeneity order, is concave in  $\mathcal{K}^n$ .

# Inequalities of Brunn-Minkowski type

# Inequalities of Brunn-Minkowski type

Let us consider a functional  $F : \mathcal{K}^n \rightarrow \mathbb{R}$  such that:

# Inequalities of Brunn-Minkowski type

Let us consider a functional  $F : \mathcal{K}^n \rightarrow \mathbb{R}$  such that:

- ▶  $F(K) \geq 0$  for every  $K \in \mathcal{K}^n$ ;

# Inequalities of Brunn-Minkowski type

Let us consider a functional  $F : \mathcal{K}^n \rightarrow \mathbb{R}$  such that:

- ▶  $F(K) \geq 0$  for every  $K \in \mathcal{K}^n$ ;
- ▶  $F$  is  $\alpha$ -homogeneous ( $\alpha \neq 0$ ):

$$F(sK) = s^\alpha F(K), \quad \forall s \geq 0, K \in \mathcal{K}^n.$$



# Inequalities of Brunn-Minkowski type

Let us consider a functional  $F : \mathcal{K}^n \rightarrow \mathbb{R}$  such that:

- ▶  $F(K) \geq 0$  for every  $K \in \mathcal{K}^n$ ;
- ▶  $F$  is  $\alpha$ -homogeneous ( $\alpha \neq 0$ ):

$$F(sK) = s^\alpha F(K), \quad \forall s \geq 0, K \in \mathcal{K}^n.$$

---

We say that  $F$  verifies a Brunn-Minkowski type inequality

# Inequalities of Brunn-Minkowski type

Let us consider a functional  $F : \mathcal{K}^n \rightarrow \mathbb{R}$  such that:

- ▶  $F(K) \geq 0$  for every  $K \in \mathcal{K}^n$ ;
- ▶  $F$  is  $\alpha$ -homogeneous ( $\alpha \neq 0$ ):

$$F(sK) = s^\alpha F(K), \quad \forall s \geq 0, K \in \mathcal{K}^n.$$

---

We say that  $F$  verifies a Brunn-Minkowski type inequality if for every  $K_0, K_1 \in \mathcal{K}^n$ , and for every  $t \in [0, 1]$ :

$$F((1-t)K_0 + tK_1)^{1/\alpha} \geq (1-t)F(K_0)^{1/\alpha} + tF(K_1)^{1/\alpha}.$$

# Examples from mathematical physics

# Examples from mathematical physics

The following functionals verify a Brunn-Minkowski type inequality.

# Examples from mathematical physics

The following functionals verify a Brunn-Minkowski type inequality.

- ▶ The **electrostatic capacity** (Borell, 1983; Caffarelli, Jerison & Lieb, 1996).

# Examples from mathematical physics

The following functionals verify a Brunn-Minkowski type inequality.

- ▶ The **electrostatic capacity** (Borell, 1983; Caffarelli, Jerison & Lieb, 1996).
- ▶ The **torsional rigidity** (Borell, 1985).

# Examples from mathematical physics

The following functionals verify a Brunn-Minkowski type inequality.

- ▶ The **electrostatic capacity** (Borell, 1983; Caffarelli, Jerison & Lieb, 1996).
- ▶ The **torsional rigidity** (Borell, 1985).
- ▶ The **principal frequency**, or first Dirichlet eigenvalue of the Laplace operator (Brascamp & Lieb, 1976).

# Examples from mathematical physics

The following functionals verify a Brunn-Minkowski type inequality.

- ▶ The **electrostatic capacity** (Borell, 1983; Caffarelli, Jerison & Lieb, 1996).
- ▶ The **torsional rigidity** (Borell, 1985).
- ▶ The **principal frequency**, or first Dirichlet eigenvalue of the Laplace operator (Brascamp & Lieb, 1976).

---

Our goal was to extend these Brunn-Minkowski inequalities to the Gauss space.



# Examples from mathematical physics

The following functionals verify a Brunn-Minkowski type inequality.

- ▶ The **electrostatic capacity** (Borell, 1983; Caffarelli, Jerison & Lieb, 1996).
- ▶ The **torsional rigidity** (Borell, 1985).
- ▶ The **principal frequency**, or first Dirichlet eigenvalue of the Laplace operator (Brascamp & Lieb, 1976).

---

Our goal was to extend these Brunn-Minkowski inequalities to the Gauss space. So far, we managed to find some new results in this direction for one of the above functionals, namely the **principal frequency**,

# Examples from mathematical physics

The following functionals verify a Brunn-Minkowski type inequality.

- ▶ The **electrostatic capacity** (Borell, 1983; Caffarelli, Jerison & Lieb, 1996).
- ▶ The **torsional rigidity** (Borell, 1985).
- ▶ The **principal frequency**, or first Dirichlet eigenvalue of the Laplace operator (Brascamp & Lieb, 1976).

---

Our goal was to extend these Brunn-Minkowski inequalities to the Gauss space. So far, we managed to find some new results in this direction for one of the above functionals, namely the **principal frequency**, on which we will now focus.

# The principal frequency of a convex body

# The principal frequency of a convex body

Let  $K \in \mathcal{K}^n$  and let  $\text{int}(K)$  be its interior;

# The principal frequency of a convex body

Let  $K \in \mathcal{K}^n$  and let  $\text{int}(K)$  be its interior; given  $v \in C^1(\text{int}(K)) \cap C(K)$ , we define its **Rayleigh quotient**:

# The principal frequency of a convex body

Let  $K \in \mathcal{K}^n$  and let  $\text{int}(K)$  be its interior; given  $v \in C^1(\text{int}(K)) \cap C(K)$ , we define its **Rayleigh quotient**:

$$R[v] = \frac{\int_K |\nabla v|^2 dx}{\int_K v^2 dx}.$$

# The principal frequency of a convex body

Let  $K \in \mathcal{K}^n$  and let  $\text{int}(K)$  be its interior; given  $v \in C^1(\text{int}(K)) \cap C(K)$ , we define its **Rayleigh quotient**:

$$R[v] = \frac{\int_K |\nabla v|^2 dx}{\int_K v^2 dx}.$$

The **principal frequency** of  $K$  is

# The principal frequency of a convex body

Let  $K \in \mathcal{K}^n$  and let  $\text{int}(K)$  be its interior; given  $v \in C^1(\text{int}(K)) \cap C(K)$ , we define its **Rayleigh quotient**:

$$R[v] = \frac{\int_K |\nabla v|^2 dx}{\int_K v^2 dx}.$$

The **principal frequency** of  $K$  is

$$\lambda(K) = \inf\{R[v]:$$



# The principal frequency of a convex body

Let  $K \in \mathcal{K}^n$  and let  $\text{int}(K)$  be its interior; given  $v \in C^1(\text{int}(K)) \cap C(K)$ , we define its **Rayleigh quotient**:

$$R[v] = \frac{\int_K |\nabla v|^2 dx}{\int_K v^2 dx}.$$

The **principal frequency** of  $K$  is

$$\lambda(K) = \inf\{R[v] : v \in C^1(\text{int}(K)) \cap C(K),$$

# The principal frequency of a convex body

Let  $K \in \mathcal{K}^n$  and let  $\text{int}(K)$  be its interior; given  $v \in C^1(\text{int}(K)) \cap C(K)$ , we define its **Rayleigh quotient**:

$$R[v] = \frac{\int_K |\nabla v|^2 dx}{\int_K v^2 dx}.$$

The **principal frequency** of  $K$  is

$$\lambda(K) = \inf\{R[v] : v \in C^1(\text{int}(K)) \cap C(K), v = 0 \text{ on } \partial K\}.$$

# The principal frequency of a convex body

Let  $K \in \mathcal{K}^n$  and let  $\text{int}(K)$  be its interior; given  $v \in C^1(\text{int}(K)) \cap C(K)$ , we define its **Rayleigh quotient**:

$$R[v] = \frac{\int_K |\nabla v|^2 dx}{\int_K v^2 dx}.$$

The **principal frequency** of  $K$  is

$$\lambda(K) = \inf\{R[v] : v \in C^1(\text{int}(K)) \cap C(K), v = 0 \text{ on } \partial K\}.$$

By its definition,  $\lambda(K)$  is also the best (i.e. the largest) possible constant  $c$  for which the following **Poincaré inequality** holds true:

# The principal frequency of a convex body

Let  $K \in \mathcal{K}^n$  and let  $\text{int}(K)$  be its interior; given  $v \in C^1(\text{int}(K)) \cap C(K)$ , we define its **Rayleigh quotient**:

$$R[v] = \frac{\int_K |\nabla v|^2 dx}{\int_K v^2 dx}.$$

The **principal frequency** of  $K$  is

$$\lambda(K) = \inf\{R[v] : v \in C^1(\text{int}(K)) \cap C(K), v = 0 \text{ on } \partial K\}.$$

By its definition,  $\lambda(K)$  is also the best (i.e. the largest) possible constant  $c$  for which the following **Poincaré inequality** holds true:

$$c \int_K v^2 dx \leq \int_K |\nabla v|^2 dx,$$

# The principal frequency of a convex body

Let  $K \in \mathcal{K}^n$  and let  $\text{int}(K)$  be its interior; given  $v \in C^1(\text{int}(K)) \cap C(K)$ , we define its **Rayleigh quotient**:

$$R[v] = \frac{\int_K |\nabla v|^2 dx}{\int_K v^2 dx}.$$

The **principal frequency** of  $K$  is

$$\lambda(K) = \inf\{R[v] : v \in C^1(\text{int}(K)) \cap C(K), v = 0 \text{ on } \partial K\}.$$

By its definition,  $\lambda(K)$  is also the best (i.e. the largest) possible constant  $c$  for which the following **Poincaré inequality** holds true:

$$c \int_K v^2 dx \leq \int_K |\nabla v|^2 dx, \quad \forall v = 0 \text{ on } \partial K.$$

$\lambda$  = first eigenvalue of the Laplace operator

$\lambda$  = first eigenvalue of the Laplace operator

The minimizer  $\bar{v}$  of the problem

$$\inf_{v|_{\partial K}=0} \frac{\int_K |\nabla v|^2 dx}{\int_K v^2 dx} = \lambda(K)$$

$\lambda$  = first eigenvalue of the Laplace operator

The minimizer  $\bar{v}$  of the problem

$$\inf_{v|_{\partial K}=0} \frac{\int_K |\nabla v|^2 dx}{\int_K v^2 dx} = \lambda(K)$$

verifies:

$$\left\{ \begin{array}{l} \Delta \bar{v} = -\lambda(K) \bar{v}, \quad \bar{v} > 0 \quad \text{in int}(K), \\ \end{array} \right.$$



$\lambda$  = first eigenvalue of the Laplace operator

The minimizer  $\bar{v}$  of the problem

$$\inf_{v|_{\partial K}=0} \frac{\int_K |\nabla v|^2 dx}{\int_K v^2 dx} = \lambda(K)$$

verifies:

$$\begin{cases} \Delta \bar{v} = -\lambda(K) \bar{v}, & \bar{v} > 0 & \text{in int}(K), \\ \bar{v} = 0 & & \text{on } \partial K. \end{cases}$$

$\lambda$  = first eigenvalue of the Laplace operator

The minimizer  $\bar{v}$  of the problem

$$\inf_{v|_{\partial K}=0} \frac{\int_K |\nabla v|^2 dx}{\int_K v^2 dx} = \lambda(K)$$

verifies:

$$\begin{cases} \Delta \bar{v} = -\lambda(K) \bar{v}, & \bar{v} > 0 & \text{in int}(K), \\ \bar{v} = 0 & & \text{on } \partial K. \end{cases}$$

Here

$$\Delta \bar{v} = \text{trace}(D^2 \bar{v}) = \sum_{i=1}^n \bar{v}_{ii},$$

is the **Laplace operator**.

# Properties of $\lambda$

# Properties of $\lambda$

$\lambda = \lambda(K)$  is:

# Properties of $\lambda$

$\lambda = \lambda(K)$  is:

- ▶ **continuous** with respect to the Hausdorff metric;

# Properties of $\lambda$

$\lambda = \lambda(K)$  is:

- ▶ **continuous** with respect to the Hausdorff metric;
- ▶ is **rigid motion invariant**;

# Properties of $\lambda$

$\lambda = \lambda(K)$  is:

- ▶ **continuous** with respect to the Hausdorff metric;
- ▶ is **rigid motion invariant**;
- ▶ **homogeneous** of degree  $-2$ :

# Properties of $\lambda$

$\lambda = \lambda(K)$  is:

- ▶ **continuous** with respect to the Hausdorff metric;
- ▶ is **rigid motion invariant**;
- ▶ **homogeneous** of degree  $-2$ :

$$\lambda(sK) = \frac{1}{s^2} \lambda(K), \quad \forall K \in \mathcal{K}^n, \forall s > 0.$$



## Detour: an isoperimetric type inequality for $\lambda$

## Detour: an isoperimetric type inequality for $\lambda$

**Theorem (Faber-Krahn inequality).** *Among all convex bodies with **fixed volume**,  $\lambda$  is minimized by balls:*

## Detour: an isoperimetric type inequality for $\lambda$

**Theorem (Faber-Krahn inequality).** *Among all convex bodies with **fixed volume**,  $\lambda$  is minimized by balls: for every  $K \in \mathcal{K}^n$ ,*

$$\lambda(K) \geq \lambda(B),$$

*where  $B$  is a ball such that  $V_n(B) = V_n(K)$ .*

## Detour: an isoperimetric type inequality for $\lambda$

**Theorem (Faber-Krahn inequality).** *Among all convex bodies with **fixed volume**,  $\lambda$  is minimized by balls: for every  $K \in \mathcal{K}^n$ ,*

$$\lambda(K) \geq \lambda(B),$$

*where  $B$  is a ball such that  $V_n(B) = V_n(K)$ .*

Behind this result there is the **Polya-Szegö principle**,

## Detour: an isoperimetric type inequality for $\lambda$

**Theorem (Faber-Krahn inequality).** *Among all convex bodies with **fixed volume**,  $\lambda$  is minimized by balls: for every  $K \in \mathcal{K}^n$ ,*

$$\lambda(K) \geq \lambda(B),$$

*where  $B$  is a ball such that  $V_n(B) = V_n(K)$ .*

Behind this result there is the **Polya-Szegö principle**, which implies in particular that the Rayleigh quotient **is monotone decreasing with respect to Steiner symmetrizations**.

# The Brunn-Minkowski inequality for $\lambda$

# The Brunn-Minkowski inequality for $\lambda$

**Theorem (Brascamp & Lieb, 1976).** *Let  $K_0, K_1 \in \mathcal{K}^n$ , and  $t \in [0, 1]$ .*

# The Brunn-Minkowski inequality for $\lambda$

**Theorem (Brascamp & Lieb, 1976).** *Let  $K_0, K_1 \in \mathcal{K}^n$ , and  $t \in [0, 1]$ . Then*

$$\lambda((1-t)K_0 + tK_1) \leq (1-t)\lambda(K_0) + t\lambda(K_1).$$



# The Brunn-Minkowski inequality for $\lambda$

**Theorem (Brascamp & Lieb, 1976).** *Let  $K_0, K_1 \in \mathcal{K}^n$ , and  $t \in [0, 1]$ . Then*

$$\lambda((1-t)K_0 + tK_1) \leq (1-t)\lambda(K_0) + t\lambda(K_1).$$

By a standard homogeneity argument, the previous result implies

$$\lambda((1-t)K_0 + tK_1)^{-1/2} \geq (1-t)\lambda(K_0)^{-1/2} + t\lambda(K_1)^{-1/2},$$

# The Brunn-Minkowski inequality for $\lambda$

**Theorem (Brascamp & Lieb, 1976).** *Let  $K_0, K_1 \in \mathcal{K}^n$ , and  $t \in [0, 1]$ . Then*

$$\lambda((1-t)K_0 + tK_1) \leq (1-t)\lambda(K_0) + t\lambda(K_1).$$

By a standard homogeneity argument, the previous result implies

$$\lambda((1-t)K_0 + tK_1)^{-1/2} \geq (1-t)\lambda(K_0)^{-1/2} + t\lambda(K_1)^{-1/2},$$

i.e. the Brunn-Minkowski inequality for  $\lambda$ .

# Back to the volume - The passage to the Gauss space

## Back to the volume - The passage to the Gauss space

We denote by  $\gamma_n$  the Gauss measure in  $\mathbb{R}^n$ :

$$\gamma_n(A) = \frac{1}{(2\pi)^{n/2}} \int_A e^{-\frac{|x|^2}{2}} dx, \quad A \subset \mathbb{R}^n, \text{ measurable.}$$

## Back to the volume - The passage to the Gauss space

We denote by  $\gamma_n$  the Gauss measure in  $\mathbb{R}^n$ :

$$\gamma_n(A) = \frac{1}{(2\pi)^{n/2}} \int_A e^{-\frac{|x|^2}{2}} dx, \quad A \subset \mathbb{R}^n, \text{ measurable.}$$

The following inequalities hold,  $\forall K_0, K_1 \in \mathcal{K}^n, \forall t \in [0, 1]$ .

## Back to the volume - The passage to the Gauss space

We denote by  $\gamma_n$  the Gauss measure in  $\mathbb{R}^n$ :

$$\gamma_n(A) = \frac{1}{(2\pi)^{n/2}} \int_A e^{-\frac{|x|^2}{2}} dx, \quad A \subset \mathbb{R}^n, \text{ measurable.}$$

The following inequalities hold,  $\forall K_0, K_1 \in \mathcal{K}^n, \forall t \in [0, 1]$ .

► The *Gaussian multiplicative Brunn-Minkowski inequality*:

$$\gamma_n((1-t)K_0 + tK_1) \geq \gamma_n(K_0)^{1-t} \gamma_n(K_1)^t$$

## Back to the volume - The passage to the Gauss space

We denote by  $\gamma_n$  the Gauss measure in  $\mathbb{R}^n$ :

$$\gamma_n(A) = \frac{1}{(2\pi)^{n/2}} \int_A e^{-\frac{|x|^2}{2}} dx, \quad A \subset \mathbb{R}^n, \text{ measurable.}$$

The following inequalities hold,  $\forall K_0, K_1 \in \mathcal{K}^n, \forall t \in [0, 1]$ .

► The *Gaussian multiplicative Brunn-Minkowski inequality*:

$$\gamma_n((1-t)K_0 + tK_1) \geq \gamma_n(K_0)^{1-t} \gamma_n(K_1)^t$$

(consequence of the Prékopa-Leindler inequality, 1971).

# Back to the volume - The passage to the Gauss space

We denote by  $\gamma_n$  the Gauss measure in  $\mathbb{R}^n$ :

$$\gamma_n(A) = \frac{1}{(2\pi)^{n/2}} \int_A e^{-\frac{|x|^2}{2}} dx, \quad A \subset \mathbb{R}^n, \text{ measurable.}$$

The following inequalities hold,  $\forall K_0, K_1 \in \mathcal{K}^n, \forall t \in [0, 1]$ .

- ▶ The *Gaussian multiplicative Brunn-Minkowski inequality*:

$$\gamma_n((1-t)K_0 + tK_1) \geq \gamma_n(K_0)^{1-t} \gamma_n(K_1)^t$$

(consequence of the Prékopa-Leindler inequality, 1971).

- ▶ The **Ehrhard inequality** (Ehrhard, 1983):

$$\Phi^{-1}(\gamma_n((1-t)K_0 + tK_1)) \geq (1-t)\Phi^{-1}(\gamma_n(K_0)) + t\Phi^{-1}(\gamma_n(K_1)),$$



# Back to the volume - The passage to the Gauss space

We denote by  $\gamma_n$  the Gauss measure in  $\mathbb{R}^n$ :

$$\gamma_n(A) = \frac{1}{(2\pi)^{n/2}} \int_A e^{-\frac{|x|^2}{2}} dx, \quad A \subset \mathbb{R}^n, \text{ measurable.}$$

The following inequalities hold,  $\forall K_0, K_1 \in \mathcal{K}^n, \forall t \in [0, 1]$ .

- ▶ The *Gaussian multiplicative Brunn-Minkowski inequality*:

$$\gamma_n((1-t)K_0 + tK_1) \geq \gamma_n(K_0)^{1-t} \gamma_n(K_1)^t$$

(consequence of the Prékopa-Leindler inequality, 1971).

- ▶ The **Ehrhard inequality** (Ehrhard, 1983):

$$\Phi^{-1}(\gamma_n((1-t)K_0 + tK_1)) \geq (1-t)\Phi^{-1}(\gamma_n(K_0)) + t\Phi^{-1}(\gamma_n(K_1)),$$

where

$$\Phi(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^x e^{-s^2/2} ds.$$

## Second detour: the isoperimetric inequality in Gauss space

## Second detour: the isoperimetric inequality in Gauss space

The Ehrhard inequality can be used to prove the Gaussian isoperimetric inequality (which is stated here for convex bodies).

## Second detour: the isoperimetric inequality in Gauss space

The Ehrhard inequality can be used to prove the Gaussian isoperimetric inequality (which is stated here for convex bodies).

**Theorem (Borell, '71, Sudakov & Tsirelson, '75).** *Among all convex bodies with fixed Gaussian measure, the half spaces minimizes the Gaussian perimeter.*

# The Gardner-Zvavitch question

# The Gardner-Zvavitch question

**Question (Gardner & Zvavitch, 2010).** Given  $K_0, K_1 \in \mathcal{K}^n$ , containing the origin, and  $t \in [0, 1]$ , is it true that

$$\gamma_n((1-t)K_0 + tK_1)^{1/n} \geq (1-t)\gamma_n(K_0)^{1/n} + t\gamma_n(K_1)^{1/n} ? \quad (*)$$

# The Gardner-Zvavitch question

**Question (Gardner & Zvavitch, 2010).** Given  $K_0, K_1 \in \mathcal{K}^n$ , containing the origin, and  $t \in [0, 1]$ , is it true that

$$\gamma_n((1-t)K_0 + tK_1)^{1/n} \geq (1-t)\gamma_n(K_0)^{1/n} + t\gamma_n(K_1)^{1/n} ? \quad (*)$$

(Dimensional Brunn-Minkowski inequality in Gauss space).

# The Gardner-Zvavitch question

**Question (Gardner & Zvavitch, 2010).** Given  $K_0, K_1 \in \mathcal{K}^n$ , containing the origin, and  $t \in [0, 1]$ , is it true that

$$\gamma_n((1-t)K_0 + tK_1)^{1/n} \geq (1-t)\gamma_n(K_0)^{1/n} + t\gamma_n(K_1)^{1/n} ? \quad (*)$$

(Dimensional Brunn-Minkowski inequality in Gauss space).

- The assumption that  $0 \in K_0, K_1$  is necessary.



# The Gardner-Zvavitch question

**Question (Gardner & Zvavitch, 2010).** Given  $K_0, K_1 \in \mathcal{K}^n$ , containing the origin, and  $t \in [0, 1]$ , is it true that

$$\gamma_n((1-t)K_0 + tK_1)^{1/n} \geq (1-t)\gamma_n(K_0)^{1/n} + t\gamma_n(K_1)^{1/n} ? \quad (*)$$

(Dimensional Brunn-Minkowski inequality in Gauss space).

- ▶ The assumption that  $0 \in K_0, K_1$  is necessary.
- ▶ (\*) is stronger than the Gaussian multiplicative Brunn-Minkowski inequality, and it is not implied by the Ehrhard inequality.

# The Gardner-Zvavitch question

**Question (Gardner & Zvavitch, 2010).** Given  $K_0, K_1 \in \mathcal{K}^n$ , containing the origin, and  $t \in [0, 1]$ , is it true that

$$\gamma_n((1-t)K_0 + tK_1)^{1/n} \geq (1-t)\gamma_n(K_0)^{1/n} + t\gamma_n(K_1)^{1/n} ? \quad (*)$$

(Dimensional Brunn-Minkowski inequality in Gauss space).

- ▶ The assumption that  $0 \in K_0, K_1$  is necessary.
- ▶ (\*) is stronger than the Gaussian multiplicative Brunn-Minkowski inequality, and it is not implied by the Ehrhard inequality.
- ▶ The exponent  $1/n$  is the best possible.

# Main steps in the solution of the Gardner-Zvavitch question

# Main steps in the solution of the Gardner-Zvavitch question

$$\gamma_n((1-t)K_0 + tK_1)^{1/n} \geq (1-t)\gamma_n(K_0)^{1/n} + t\gamma_n(K_1)^{1/n}$$

# Main steps in the solution of the Gardner-Zvavitch question

$$\gamma_n((1-t)K_0 + tK_1)^{1/n} \geq (1-t)\gamma_n(K_0)^{1/n} + t\gamma_n(K_1)^{1/n}$$

- ▶ Gardner & Zvavitch (2010) verified that the answer is “yes” for coordinate boxes and slabs, containing the origin, and dilatates of a common origin-symmetric convex body

# Main steps in the solution of the Gardner-Zvavitch question

$$\gamma_n((1-t)K_0 + tK_1)^{1/n} \geq (1-t)\gamma_n(K_0)^{1/n} + t\gamma_n(K_1)^{1/n}$$

- ▶ Gardner & Zvavitch (2010) verified that the answer is “yes” for coordinate boxes and slabs, containing the origin, and dilatates of a common origin-symmetric convex body (this last case shows the connection of their question with the B-conjecture).

# Main steps in the solution of the Gardner-Zvavitch question

$$\gamma_n((1-t)K_0 + tK_1)^{1/n} \geq (1-t)\gamma_n(K_0)^{1/n} + t\gamma_n(K_1)^{1/n}$$

- ▶ Gardner & Zvavitch (2010) verified that the answer is “yes” for coordinate boxes and slabs, containing the origin, and dilatates of a common origin-symmetric convex body (this last case shows the connection of their question with the B-conjecture).
- ▶ Nayar & Tkocz (2013) showed by counterexamples that the answer is “no” for general convex bodies containing the origin.

# Main steps in the solution of the Gardner-Zvavitch question

$$\gamma_n((1-t)K_0 + tK_1)^{1/n} \geq (1-t)\gamma_n(K_0)^{1/n} + t\gamma_n(K_1)^{1/n}$$

- ▶ Gardner & Zvavitch (2010) verified that the answer is “yes” for coordinate boxes and slabs, containing the origin, and dilatates of a common origin-symmetric convex body (this last case shows the connection of their question with the B-conjecture).
- ▶ Nayar & Tkocz (2013) showed by counterexamples that the answer is “no” for general convex bodies containing the origin.
- ▶ Kolesnikov & Livshyts (2021) proved the inequality for origin symmetric convex bodies with the exponent  $1/n$  replaced by  $2/n$ .



# Main steps in the solution of the Gardner-Zvavitch question

$$\gamma_n((1-t)K_0 + tK_1)^{1/n} \geq (1-t)\gamma_n(K_0)^{1/n} + t\gamma_n(K_1)^{1/n}$$

- ▶ Gardner & Zvavitch (2010) verified that the answer is “yes” for coordinate boxes and slabs, containing the origin, and dilatates of a common origin-symmetric convex body (this last case shows the connection of their question with the B-conjecture).
- ▶ Nayar & Tkocz (2013) showed by counterexamples that the answer is “no” for general convex bodies containing the origin.
- ▶ Kolesnikov & Livshyts (2021) proved the inequality for origin symmetric convex bodies with the exponent  $1/n$  replaced by  $2/n$ .
- ▶ The affirmative answer to the question for origin-symmetric convex bodies was given by Eskenazis & Moschidis (2021).

# The Gaussian principal frequency

# The Gaussian principal frequency

Let  $K \in \mathcal{K}^n$ ; we set, for  $v \in C^1(\text{int}(K)) \cap C(K)$ ,

$$R_\gamma[v] = \frac{\int_K |\nabla v|^2 d\gamma_n}{\int_K v^2 d\gamma_n},$$

# The Gaussian principal frequency

Let  $K \in \mathcal{K}^n$ ; we set, for  $v \in C^1(\text{int}(K)) \cap C(K)$ ,

$$R_\gamma[v] = \frac{\int_K |\nabla v|^2 d\gamma_n}{\int_K v^2 d\gamma_n},$$

and

$$\lambda_\gamma(K)$$

# The Gaussian principal frequency

Let  $K \in \mathcal{K}^n$ ; we set, for  $v \in C^1(\text{int}(K)) \cap C(K)$ ,

$$R_\gamma[v] = \frac{\int_K |\nabla v|^2 d\gamma_n}{\int_K v^2 d\gamma_n},$$

and

$$\lambda_\gamma(K) = \inf\{R_\gamma[v]:$$

# The Gaussian principal frequency

Let  $K \in \mathcal{K}^n$ ; we set, for  $v \in C^1(\text{int}(K)) \cap C(K)$ ,

$$R_\gamma[v] = \frac{\int_K |\nabla v|^2 d\gamma_n}{\int_K v^2 d\gamma_n},$$

and

$$\lambda_\gamma(K) = \inf\{R_\gamma[v] : v \in C^1(\text{int}(K)) \cap C(K),$$

# The Gaussian principal frequency

Let  $K \in \mathcal{K}^n$ ; we set, for  $v \in C^1(\text{int}(K)) \cap C(K)$ ,

$$R_\gamma[v] = \frac{\int_K |\nabla v|^2 d\gamma_n}{\int_K v^2 d\gamma_n},$$

and

$$\lambda_\gamma(K) = \inf\{R_\gamma[v] : v \in C^1(\text{int}(K)) \cap C(K), v = 0 \text{ on } \partial K\}.$$

# The Gaussian principal frequency

Let  $K \in \mathcal{K}^n$ ; we set, for  $v \in C^1(\text{int}(K)) \cap C(K)$ ,

$$R_\gamma[v] = \frac{\int_K |\nabla v|^2 d\gamma_n}{\int_K v^2 d\gamma_n},$$

and

$$\lambda_\gamma(K) = \inf\{R_\gamma[v] : v \in C^1(\text{int}(K)) \cap C(K), v = 0 \text{ on } \partial K\}.$$

- $\lambda_\gamma(K)$  is the largest possible constant  $c$  for which the following **Gaussian Poincaré inequality** holds true:



# The Gaussian principal frequency

Let  $K \in \mathcal{K}^n$ ; we set, for  $v \in C^1(\text{int}(K)) \cap C(K)$ ,

$$R_\gamma[v] = \frac{\int_K |\nabla v|^2 d\gamma_n}{\int_K v^2 d\gamma_n},$$

and

$$\lambda_\gamma(K) = \inf\{R_\gamma[v] : v \in C^1(\text{int}(K)) \cap C(K), v = 0 \text{ on } \partial K\}.$$

- ▶  $\lambda_\gamma(K)$  is the largest possible constant  $c$  for which the following **Gaussian Poincaré inequality** holds true:

$$c \int_K v^2 d\gamma_n \leq \int_K |\nabla v|^2 d\gamma_n$$

# The Gaussian principal frequency

Let  $K \in \mathcal{K}^n$ ; we set, for  $v \in C^1(\text{int}(K)) \cap C(K)$ ,

$$R_\gamma[v] = \frac{\int_K |\nabla v|^2 d\gamma_n}{\int_K v^2 d\gamma_n},$$

and

$$\lambda_\gamma(K) = \inf\{R_\gamma[v] : v \in C^1(\text{int}(K)) \cap C(K), v = 0 \text{ on } \partial K\}.$$

- $\lambda_\gamma(K)$  is the largest possible constant  $c$  for which the following **Gaussian Poincaré inequality** holds true:

$$c \int_K v^2 d\gamma_n \leq \int_K |\nabla v|^2 d\gamma_n, \quad \forall v = 0 \text{ on } \partial K.$$

# The Gaussian principal frequency

Let  $K \in \mathcal{K}^n$ ; we set, for  $v \in C^1(\text{int}(K)) \cap C(K)$ ,

$$R_\gamma[v] = \frac{\int_K |\nabla v|^2 d\gamma_n}{\int_K v^2 d\gamma_n},$$

and

$$\lambda_\gamma(K) = \inf\{R_\gamma[v] : v \in C^1(\text{int}(K)) \cap C(K), v = 0 \text{ on } \partial K\}.$$

- ▶  $\lambda_\gamma(K)$  is the largest possible constant  $c$  for which the following **Gaussian Poincaré inequality** holds true:

$$c \int_K v^2 d\gamma_n \leq \int_K |\nabla v|^2 d\gamma_n, \quad \forall v = 0 \text{ on } \partial K.$$

- ▶  $\lambda_\gamma$  is rotation invariant, but it is **neither translation invariant, nor homogeneous**.

# The associated boundary value problem

# The associated boundary value problem

The minimiser  $\bar{v}$  of the problem

$$\inf_{v|_{\partial K}=0} \frac{\int_K |\nabla v|^2 d\gamma_n}{\int_K v^2 d\gamma_n} = \lambda_\gamma(K)$$

# The associated boundary value problem

The minimiser  $\bar{v}$  of the problem

$$\inf_{v|_{\partial K}=0} \frac{\int_K |\nabla v|^2 d\gamma_n}{\int_K v^2 d\gamma_n} = \lambda_\gamma(K)$$

verifies:

$$\begin{cases} \Delta \bar{v} - \langle \nabla \bar{v}, x \rangle = -\lambda_\gamma(K) \bar{v}, & \bar{v} > 0 & \text{in int}(K), \end{cases}$$

# The associated boundary value problem

The minimiser  $\bar{v}$  of the problem

$$\inf_{v|_{\partial K}=0} \frac{\int_K |\nabla v|^2 d\gamma_n}{\int_K v^2 d\gamma_n} = \lambda_\gamma(K)$$

verifies:

$$\begin{cases} \Delta \bar{v} - \langle \nabla \bar{v}, x \rangle = -\lambda_\gamma(K) \bar{v}, & \bar{v} > 0 & \text{in int}(K), \\ \bar{v} = 0 & & \text{on } \partial K. \end{cases}$$

# The associated boundary value problem

The minimiser  $\bar{v}$  of the problem

$$\inf_{v|_{\partial K}=0} \frac{\int_K |\nabla v|^2 d\gamma_n}{\int_K v^2 d\gamma_n} = \lambda_\gamma(K)$$

verifies:

$$\begin{cases} \Delta \bar{v} - \langle \nabla \bar{v}, x \rangle = -\lambda_\gamma(K) \bar{v}, & \bar{v} > 0 & \text{in int}(K), \\ \bar{v} = 0 & & \text{on } \partial K. \end{cases}$$

$$u \mapsto \Delta u - \langle \nabla u, x \rangle$$



# The associated boundary value problem

The minimiser  $\bar{v}$  of the problem

$$\inf_{v|_{\partial K}=0} \frac{\int_K |\nabla v|^2 d\gamma_n}{\int_K v^2 d\gamma_n} = \lambda_\gamma(K)$$

verifies:

$$\begin{cases} \Delta \bar{v} - \langle \nabla \bar{v}, x \rangle = -\lambda_\gamma(K) \bar{v}, & \bar{v} > 0 & \text{in int}(K), \\ \bar{v} = 0 & & \text{on } \partial K. \end{cases}$$

$$u \mapsto \Delta u - \langle \nabla u, x \rangle$$

is the Hermite, or Ornstein-Uhlenbeck, operator.

Third detour: an isoperimetric type inequality for  $\lambda_\gamma$

## Third detour: an isoperimetric type inequality for $\lambda_\gamma$

**Theorem (Ehrhard, 1984).** *Among all convex bodies with fixed Gaussian volume,*

## Third detour: an isoperimetric type inequality for $\lambda_\gamma$

**Theorem (Ehrhard, 1984).** *Among all convex bodies with fixed Gaussian volume,  $\lambda_\gamma$  is minimized by **half-spaces**:*

## Third detour: an isoperimetric type inequality for $\lambda_\gamma$

**Theorem (Ehrhard, 1984).** *Among all convex bodies with fixed Gaussian volume,  $\lambda_\gamma$  is minimized by **half-spaces**: for every  $K \in \mathcal{K}^n$ ,*

$$\lambda_\gamma(K) \geq \lambda_\gamma(H)$$

*where  $H$  is any half-space such that  $\gamma_n(K) = \gamma_n(H)$ .*

## Third detour: an isoperimetric type inequality for $\lambda_\gamma$

**Theorem (Ehrhard, 1984).** *Among all convex bodies with fixed Gaussian volume,  $\lambda_\gamma$  is minimized by **half-spaces**: for every  $K \in \mathcal{K}^n$ ,*

$$\lambda_\gamma(K) \geq \lambda_\gamma(H)$$

*where  $H$  is any half-space such that  $\gamma_n(K) = \gamma_n(H)$ .*

- ▶ This result fully parallels the Gaussian isoperimetric inequality.

## Third detour: an isoperimetric type inequality for $\lambda_\gamma$

**Theorem (Ehrhard, 1984).** *Among all convex bodies with fixed Gaussian volume,  $\lambda_\gamma$  is minimized by **half-spaces**: for every  $K \in \mathcal{K}^n$ ,*

$$\lambda_\gamma(K) \geq \lambda_\gamma(H)$$

*where  $H$  is any half-space such that  $\gamma_n(K) = \gamma_n(H)$ .*

- ▶ This result fully parallels the Gaussian isoperimetric inequality.
- ▶ The proof exploits a (de-)symmetrization procedure (*Gaussian symmetrization*),

## Third detour: an isoperimetric type inequality for $\lambda_\gamma$

**Theorem (Ehrhard, 1984).** *Among all convex bodies with fixed Gaussian volume,  $\lambda_\gamma$  is minimized by **half-spaces**: for every  $K \in \mathcal{K}^n$ ,*

$$\lambda_\gamma(K) \geq \lambda_\gamma(H)$$

*where  $H$  is any half-space such that  $\gamma_n(K) = \gamma_n(H)$ .*

- ▶ This result fully parallels the Gaussian isoperimetric inequality.
- ▶ The proof exploits a (de-)symmetrization procedure (*Gaussian symmetrization*), which preserves the Gaussian measure and reduces  $\lambda_\gamma$



## Third detour: an isoperimetric type inequality for $\lambda_\gamma$

**Theorem (Ehrhard, 1984).** *Among all convex bodies with fixed Gaussian volume,  $\lambda_\gamma$  is minimized by **half-spaces**: for every  $K \in \mathcal{K}^n$ ,*

$$\lambda_\gamma(K) \geq \lambda_\gamma(H)$$

*where  $H$  is any half-space such that  $\gamma_n(K) = \gamma_n(H)$ .*

- ▶ This result fully parallels the Gaussian isoperimetric inequality.
- ▶ The proof exploits a (de-)symmetrization procedure (*Gaussian symmetrization*), which preserves the Gaussian measure and reduces  $\lambda_\gamma$  (and the Gaussian perimeter).

# A convexity inequality for $\lambda_\gamma$

## A convexity inequality for $\lambda_\gamma$

**Theorem (Francini, Livshyts, Salani, C. - 2023).** *Let  $K_0, K_1 \in \mathcal{K}^n$  be origin symmetric convex bodies, and let  $t \in [0, 1]$ .*

## A convexity inequality for $\lambda_\gamma$

**Theorem (Francini, Livshyts, Salani, C. - 2023).** *Let  $K_0, K_1 \in \mathcal{K}^n$  be origin symmetric convex bodies, and let  $t \in [0, 1]$ . Then*

$$\lambda_\gamma((1-t)K_0 + tK_1) \leq (1-t)\lambda_\gamma(K_0) + t\lambda_\gamma(K_1).$$

## A convexity inequality for $\lambda_\gamma$

**Theorem (Francini, Livshyts, Salani, C. - 2023).** *Let  $K_0, K_1 \in \mathcal{K}^n$  be origin symmetric convex bodies, and let  $t \in [0, 1]$ . Then*

$$\lambda_\gamma((1-t)K_0 + tK_1) \leq (1-t)\lambda_\gamma(K_0) + t\lambda_\gamma(K_1). \quad (*)$$

- Inequality (\*) coincides with that found by Brascamp & Lieb for  $\lambda$  in the standard case (i.e with the Lebesgue measure in place of the Gaussian measure).

## A convexity inequality for $\lambda_\gamma$

**Theorem (Francini, Livshyts, Salani, C. - 2023).** *Let  $K_0, K_1 \in \mathcal{K}^n$  be origin symmetric convex bodies, and let  $t \in [0, 1]$ . Then*

$$\lambda_\gamma((1-t)K_0 + tK_1) \leq (1-t)\lambda_\gamma(K_0) + t\lambda_\gamma(K_1). \quad (*)$$

- ▶ Inequality (\*) coincides with that found by Brascamp & Lieb for  $\lambda$  in the standard case (i.e with the Lebesgue measure in place of the Gaussian measure).
- ▶ Due to the lack of homogeneity, (\*) does not imply the stronger inequality:

$$\lambda_\gamma((1-t)K_0 + tK_1)^{-1/2} \geq (1-t)\lambda_\gamma(K_0)^{-1/2} + t\lambda_\gamma(K_1)^{-1/2}.$$

# Towards the proof - A crucial tool

# Towards the proof - A crucial tool

**Theorem (FLSC - 2023).** *Let  $K \in \mathcal{K}^n$  be origin symmetric. Let  $u$  be a solution of the boundary value problem:*

$$\begin{cases} \Delta v - \langle \nabla v, x \rangle = -\lambda_\gamma(K)v, & v > 0 & \text{in } \text{int}(K), \\ v = 0 & & \text{on } \partial K. \end{cases}$$



# Towards the proof - A crucial tool

**Theorem (FLSC - 2023).** *Let  $K \in \mathcal{K}^n$  be origin symmetric. Let  $u$  be a solution of the boundary value problem:*

$$\begin{cases} \Delta v - \langle \nabla v, x \rangle = -\lambda_\gamma(K)v, & v > 0 & \text{in } \text{int}(K), \\ v = 0 & & \text{on } \partial K. \end{cases}$$

*Then  $u$  is **log-concave**.*

# Towards the proof - A crucial tool

**Theorem (FLSC - 2023).** *Let  $K \in \mathcal{K}^n$  be origin symmetric. Let  $u$  be a solution of the boundary value problem:*

$$\begin{cases} \Delta v - \langle \nabla v, x \rangle = -\lambda_\gamma(K)v, & v > 0 & \text{in } \text{int}(K), \\ v = 0 & & \text{on } \partial K. \end{cases}$$

*Then  $u$  is **log-concave**.*

The same property holds in the standard case (Brascamp & Lieb, 1976).

# Towards the proof - A crucial tool

**Theorem (FLSC - 2023).** *Let  $K \in \mathcal{K}^n$  be origin symmetric. Let  $u$  be a solution of the boundary value problem:*

$$\begin{cases} \Delta v - \langle \nabla v, x \rangle = -\lambda_\gamma(K)v, & v > 0 & \text{in } \text{int}(K), \\ v = 0 & & \text{on } \partial K. \end{cases}$$

*Then  $u$  is **log-concave**.*

The same property holds in the standard case (Brascamp & Lieb, 1976).

Several approaches, developed in the standard case, can be adapted to the Gaussian case.

## Towards the proof - A crucial tool

**Theorem (FLSC - 2023).** *Let  $K \in \mathcal{K}^n$  be origin symmetric. Let  $u$  be a solution of the boundary value problem:*

$$\begin{cases} \Delta v - \langle \nabla v, x \rangle = -\lambda_\gamma(K)v, & v > 0 & \text{in } \text{int}(K), \\ v = 0 & & \text{on } \partial K. \end{cases}$$

*Then  $u$  is **log-concave**.*

The same property holds in the standard case (Brascamp & Lieb, 1976).

Several approaches, developed in the standard case, can be adapted to the Gaussian case.

E. Milman suggested to us an additional argument based on the original proof by Brascamp & Lieb.

# The proof of the convexity inequality

## The proof of the convexity inequality

$$\lambda_\gamma((1-t)K_0 + tK_1) \leq (1-t)\lambda_\gamma(K_0) + t\lambda_\gamma(K_1)$$

## The proof of the convexity inequality

$$\lambda_\gamma((1-t)K_0 + tK_1) \leq (1-t)\lambda_\gamma(K_0) + t\lambda_\gamma(K_1)$$

Given  $K_0$  and  $K_1$  we consider  $v_0$  and  $v_1$  such that

# The proof of the convexity inequality

$$\lambda_\gamma((1-t)K_0 + tK_1) \leq (1-t)\lambda_\gamma(K_0) + t\lambda_\gamma(K_1)$$

Given  $K_0$  and  $K_1$  we consider  $v_0$  and  $v_1$  such that

$$\begin{cases} \Delta v_0 - \langle \nabla v_0, x \rangle = -\lambda_\gamma(K_0)v_0, & v_0 > 0 & \text{in } \text{int}(K_0), \\ v_0 = 0 & & \text{on } \partial K_0. \end{cases}$$



# The proof of the convexity inequality

$$\lambda_\gamma((1-t)K_0 + tK_1) \leq (1-t)\lambda_\gamma(K_0) + t\lambda_\gamma(K_1)$$

Given  $K_0$  and  $K_1$  we consider  $v_0$  and  $v_1$  such that

$$\begin{cases} \Delta v_0 - \langle \nabla v_0, x \rangle = -\lambda_\gamma(K_0)v_0, & v_0 > 0 & \text{in int}(K_0), \\ v_0 = 0 & & \text{on } \partial K_0. \end{cases}$$
$$\begin{cases} \Delta v_1 - \langle \nabla v_1, x \rangle = -\lambda_\gamma(K_1)v_1, & v_1 > 0 & \text{in int}(K_1), \\ v_1 = 0 & & \text{on } \partial K_1. \end{cases}$$

# The proof of the convexity inequality

$$\lambda_\gamma((1-t)K_0 + tK_1) \leq (1-t)\lambda_\gamma(K_0) + t\lambda_\gamma(K_1)$$

Given  $K_0$  and  $K_1$  we consider  $v_0$  and  $v_1$  such that

$$\begin{cases} \Delta v_0 - \langle \nabla v_0, x \rangle = -\lambda_\gamma(K_0)v_0, & v_0 > 0 & \text{in int}(K_0), \\ v_0 = 0 & & \text{on } \partial K_0. \end{cases}$$
$$\begin{cases} \Delta v_1 - \langle \nabla v_1, x \rangle = -\lambda_\gamma(K_1)v_1, & v_1 > 0 & \text{in int}(K_1), \\ v_1 = 0 & & \text{on } \partial K_1. \end{cases}$$

$$v_0 = e^{-u_0},$$

# The proof of the convexity inequality

$$\lambda_\gamma((1-t)K_0 + tK_1) \leq (1-t)\lambda_\gamma(K_0) + t\lambda_\gamma(K_1)$$

Given  $K_0$  and  $K_1$  we consider  $v_0$  and  $v_1$  such that

$$\begin{cases} \Delta v_0 - \langle \nabla v_0, x \rangle = -\lambda_\gamma(K_0)v_0, & v_0 > 0 & \text{in int}(K_0), \\ v_0 = 0 & & \text{on } \partial K_0. \end{cases}$$
$$\begin{cases} \Delta v_1 - \langle \nabla v_1, x \rangle = -\lambda_\gamma(K_1)v_1, & v_1 > 0 & \text{in int}(K_1), \\ v_1 = 0 & & \text{on } \partial K_1. \end{cases}$$

$$v_0 = e^{-u_0}, \quad v_1 = e^{-u_1},$$

# The proof of the convexity inequality

$$\lambda_\gamma((1-t)K_0 + tK_1) \leq (1-t)\lambda_\gamma(K_0) + t\lambda_\gamma(K_1)$$

Given  $K_0$  and  $K_1$  we consider  $v_0$  and  $v_1$  such that

$$\begin{cases} \Delta v_0 - \langle \nabla v_0, x \rangle = -\lambda_\gamma(K_0)v_0, & v_0 > 0 & \text{in int}(K_0), \\ v_0 = 0 & & \text{on } \partial K_0. \end{cases}$$
$$\begin{cases} \Delta v_1 - \langle \nabla v_1, x \rangle = -\lambda_\gamma(K_1)v_1, & v_1 > 0 & \text{in int}(K_1), \\ v_1 = 0 & & \text{on } \partial K_1. \end{cases}$$

$$v_0 = e^{-u_0}, \quad v_1 = e^{-u_1}, \quad u_0, u_1 \text{ convex.}$$

# The proof of the convexity inequality

$$\lambda_\gamma((1-t)K_0 + tK_1) \leq (1-t)\lambda_\gamma(K_0) + t\lambda_\gamma(K_1)$$

Given  $K_0$  and  $K_1$  we consider  $v_0$  and  $v_1$  such that

$$\begin{cases} \Delta v_0 - \langle \nabla v_0, x \rangle = -\lambda_\gamma(K_0)v_0, & v_0 > 0 & \text{in int}(K_0), \\ v_0 = 0 & & \text{on } \partial K_0. \end{cases}$$
$$\begin{cases} \Delta v_1 - \langle \nabla v_1, x \rangle = -\lambda_\gamma(K_1)v_1, & v_1 > 0 & \text{in int}(K_1), \\ v_1 = 0 & & \text{on } \partial K_1. \end{cases}$$

$$v_0 = e^{-u_0}, \quad v_1 = e^{-u_1}, \quad u_0, u_1 \text{ convex.}$$

We set, for  $z \in K_t = (1-t)K_0 + tK_1$ ,

# The proof of the convexity inequality

$$\lambda_\gamma((1-t)K_0 + tK_1) \leq (1-t)\lambda_\gamma(K_0) + t\lambda_\gamma(K_1)$$

Given  $K_0$  and  $K_1$  we consider  $v_0$  and  $v_1$  such that

$$\begin{cases} \Delta v_0 - \langle \nabla v_0, x \rangle = -\lambda_\gamma(K_0)v_0, & v_0 > 0 & \text{in int}(K_0), \\ v_0 = 0 & & \text{on } \partial K_0. \end{cases}$$
$$\begin{cases} \Delta v_1 - \langle \nabla v_1, x \rangle = -\lambda_\gamma(K_1)v_1, & v_1 > 0 & \text{in int}(K_1), \\ v_1 = 0 & & \text{on } \partial K_1. \end{cases}$$

$$v_0 = e^{-u_0}, \quad v_1 = e^{-u_1}, \quad u_0, u_1 \text{ convex.}$$

We set, for  $z \in K_t = (1-t)K_0 + tK_1$ ,

$$u_t(z)$$

# The proof of the convexity inequality

$$\lambda_\gamma((1-t)K_0 + tK_1) \leq (1-t)\lambda_\gamma(K_0) + t\lambda_\gamma(K_1)$$

Given  $K_0$  and  $K_1$  we consider  $v_0$  and  $v_1$  such that

$$\begin{cases} \Delta v_0 - \langle \nabla v_0, x \rangle = -\lambda_\gamma(K_0)v_0, & v_0 > 0 & \text{in int}(K_0), \\ v_0 = 0 & & \text{on } \partial K_0. \end{cases}$$
$$\begin{cases} \Delta v_1 - \langle \nabla v_1, x \rangle = -\lambda_\gamma(K_1)v_1, & v_1 > 0 & \text{in int}(K_1), \\ v_1 = 0 & & \text{on } \partial K_1. \end{cases}$$

$$v_0 = e^{-u_0}, \quad v_1 = e^{-u_1}, \quad u_0, u_1 \text{ convex.}$$

We set, for  $z \in K_t = (1-t)K_0 + tK_1$ ,

$$u_t(z) = \inf_{(1-t)x+ty=z} (1-t)u_0(x) + tu_1(y)$$

# The proof of the convexity inequality

$$\lambda_\gamma((1-t)K_0 + tK_1) \leq (1-t)\lambda_\gamma(K_0) + t\lambda_\gamma(K_1)$$

Given  $K_0$  and  $K_1$  we consider  $v_0$  and  $v_1$  such that

$$\begin{cases} \Delta v_0 - \langle \nabla v_0, x \rangle = -\lambda_\gamma(K_0)v_0, & v_0 > 0 & \text{in int}(K_0), \\ v_0 = 0 & & \text{on } \partial K_0. \end{cases}$$
$$\begin{cases} \Delta v_1 - \langle \nabla v_1, x \rangle = -\lambda_\gamma(K_1)v_1, & v_1 > 0 & \text{in int}(K_1), \\ v_1 = 0 & & \text{on } \partial K_1. \end{cases}$$

$$v_0 = e^{-u_0}, \quad v_1 = e^{-u_1}, \quad u_0, u_1 \text{ convex.}$$

We set, for  $z \in K_t = (1-t)K_0 + tK_1$ ,

$$u_t(z) = \inf_{(1-t)x+ty=z} (1-t)u_0(x) + tu_1(y) = (1-t) \cdot u_0 \square t \cdot u_1.$$



# The proof of the convexity inequality

$$\lambda_\gamma((1-t)K_0 + tK_1) \leq (1-t)\lambda_\gamma(K_0) + t\lambda_\gamma(K_1)$$

Given  $K_0$  and  $K_1$  we consider  $v_0$  and  $v_1$  such that

$$\begin{cases} \Delta v_0 - \langle \nabla v_0, x \rangle = -\lambda_\gamma(K_0)v_0, & v_0 > 0 & \text{in int}(K_0), \\ v_0 = 0 & & \text{on } \partial K_0. \end{cases}$$
$$\begin{cases} \Delta v_1 - \langle \nabla v_1, x \rangle = -\lambda_\gamma(K_1)v_1, & v_1 > 0 & \text{in int}(K_1), \\ v_1 = 0 & & \text{on } \partial K_1. \end{cases}$$

$$v_0 = e^{-u_0}, \quad v_1 = e^{-u_1}, \quad u_0, u_1 \text{ convex.}$$

We set, for  $z \in K_t = (1-t)K_0 + tK_1$ ,

$$u_t(z) = \inf_{(1-t)x+ty=z} (1-t)u_0(x) + tu_1(y) = (1-t) \cdot u_0 \square t \cdot u_1.$$

$u_t$  is convex in  $K_t$ .

# The proof of the convexity inequality

$$\lambda_\gamma((1-t)K_0 + tK_1) \leq (1-t)\lambda_\gamma(K_0) + t\lambda_\gamma(K_1)$$

Given  $K_0$  and  $K_1$  we consider  $v_0$  and  $v_1$  such that

$$\begin{cases} \Delta v_0 - \langle \nabla v_0, x \rangle = -\lambda_\gamma(K_0)v_0, & v_0 > 0 & \text{in int}(K_0), \\ v_0 = 0 & & \text{on } \partial K_0. \end{cases}$$
$$\begin{cases} \Delta v_1 - \langle \nabla v_1, x \rangle = -\lambda_\gamma(K_1)v_1, & v_1 > 0 & \text{in int}(K_1), \\ v_1 = 0 & & \text{on } \partial K_1. \end{cases}$$

$$v_0 = e^{-u_0}, \quad v_1 = e^{-u_1}, \quad u_0, u_1 \text{ convex.}$$

We set, for  $z \in K_t = (1-t)K_0 + tK_1$ ,

$$u_t(z) = \inf_{(1-t)x+ty=z} (1-t)u_0(x) + tu_1(y) = (1-t) \cdot u_0 \square t \cdot u_1.$$

$u_t$  is convex in  $K_t$ . We also set

$$v_t = e^{-u_t}.$$

$$u_t(z) = \inf_{(1-t)x+ty=z} (1-t)u_0(x) + tu_1(y)$$

$$u_t(z) = \inf_{(1-t)x+ty=z} (1-t)u_0(x) + tu_1(y)$$

For every  $z \in K_t$ ,

$$u_t(z) = \inf_{(1-t)x+ty=z} (1-t)u_0(x) + tu_1(y)$$

For every  $z \in K_t$ , there exist  $x \in K_0, y \in K_1$ , s.t.

$$z = (1-t)x + ty$$

$$u_t(z) = \inf_{(1-t)x+ty=z} (1-t)u_0(x) + tu_1(y)$$

For every  $z \in K_t$ , there exist  $x \in K_0, y \in K_1$ , s.t.

$$z = (1-t)x + ty \quad \text{and} \quad u_t(z) = (1-t)u_0(x) + tu_1(y),$$

$$u_t(z) = \inf_{(1-t)x+ty=z} (1-t)u_0(x) + tu_1(y)$$

For every  $z \in K_t$ , there exist  $x \in K_0, y \in K_1$ , s.t.

$$z = (1-t)x + ty \quad \text{and} \quad u_t(z) = (1-t)u_0(x) + tu_1(y),$$

$$\nabla u_t(z) = \nabla u_0(x) = \nabla u_1(y),$$

$$u_t(z) = \inf_{(1-t)x+ty=z} (1-t)u_0(x) + tu_1(y)$$

For every  $z \in K_t$ , there exist  $x \in K_0, y \in K_1$ , s.t.

$$z = (1-t)x + ty \quad \text{and} \quad u_t(z) = (1-t)u_0(x) + tu_1(y),$$

$$\nabla u_t(z) = \nabla u_0(x) = \nabla u_1(y),$$

$$D^2 u_t(z) = [(1-t)D^2 u_0(x)^{-1} + tD^2 u_1(y)^{-1}]^{-1}.$$



$$u_t(z) = \inf_{(1-t)x+ty=z} (1-t)u_0(x) + tu_1(y)$$

For every  $z \in K_t$ , there exist  $x \in K_0, y \in K_1$ , s.t.

$$z = (1-t)x + ty \quad \text{and} \quad u_t(z) = (1-t)u_0(x) + tu_1(y),$$

$$\nabla u_t(z) = \nabla u_0(x) = \nabla u_1(y),$$

$$D^2 u_t(z) = [(1-t)D^2 u_0(x)^{-1} + tD^2 u_1(y)^{-1}]^{-1}.$$

Using these facts,

$$u_t(z) = \inf_{(1-t)x+ty=z} (1-t)u_0(x) + tu_1(y)$$

For every  $z \in K_t$ , there exist  $x \in K_0, y \in K_1$ , s.t.

$$z = (1-t)x + ty \quad \text{and} \quad u_t(z) = (1-t)u_0(x) + tu_1(y),$$

$$\nabla u_t(z) = \nabla u_0(x) = \nabla u_1(y),$$

$$D^2 u_t(z) = [(1-t)D^2 u_0(x)^{-1} + tD^2 u_1(y)^{-1}]^{-1}.$$

Using these facts, the equations verified by  $v_0 = e^{-u_0}$  and  $v_1 = e^{-u_1}$ ,

$$u_t(z) = \inf_{(1-t)x+ty=z} (1-t)u_0(x) + tu_1(y)$$

For every  $z \in K_t$ , there exist  $x \in K_0, y \in K_1$ , s.t.

$$z = (1-t)x + ty \quad \text{and} \quad u_t(z) = (1-t)u_0(x) + tu_1(y),$$

$$\nabla u_t(z) = \nabla u_0(x) = \nabla u_1(y),$$

$$D^2 u_t(z) = [(1-t)D^2 u_1(x)^{-1} + tD^2 u_1(y)^{-1}]^{-1}.$$

Using these facts, the equations verified by  $v_0 = e^{-u_0}$  and  $v_1 = e^{-u_1}$ , the convexity of the application

$$M \mapsto \text{trace}(M^{-1}), \quad (M > 0),$$

$$u_t(z) = \inf_{(1-t)x+ty=z} (1-t)u_0(x) + tu_1(y)$$

For every  $z \in K_t$ , there exist  $x \in K_0, y \in K_1$ , s.t.

$$z = (1-t)x + ty \quad \text{and} \quad u_t(z) = (1-t)u_0(x) + tu_1(y),$$

$$\nabla u_t(z) = \nabla u_0(x) = \nabla u_1(y),$$

$$D^2 u_t(z) = [(1-t)D^2 u_0(x)^{-1} + tD^2 u_1(y)^{-1}]^{-1}.$$

Using these facts, the equations verified by  $v_0 = e^{-u_0}$  and  $v_1 = e^{-u_1}$ , the convexity of the application

$$M \mapsto \text{trace}(M^{-1}), \quad (M > 0),$$

we deduce that  $v_t = e^{-u_t}$  verifies the differential inequality:

$$u_t(z) = \inf_{(1-t)x+ty=z} (1-t)u_0(x) + tu_1(y)$$

For every  $z \in K_t$ , there exist  $x \in K_0, y \in K_1$ , s.t.

$$\begin{aligned} z &= (1-t)x + ty \quad \text{and} \quad u_t(z) = (1-t)u_0(x) + tu_1(y), \\ \nabla u_t(z) &= \nabla u_0(x) = \nabla u_1(y), \\ D^2 u_t(z) &= [(1-t)D^2 u_1(x)^{-1} + tD^2 u_1(y)^{-1}]^{-1}. \end{aligned}$$

Using these facts, the equations verified by  $v_0 = e^{-u_0}$  and  $v_1 = e^{-u_1}$ , the convexity of the application

$$M \mapsto \text{trace}(M^{-1}), \quad (M > 0),$$

we deduce that  $v_t = e^{-u_t}$  verifies the differential inequality:

$$\Delta v_t - \langle \nabla v_t, z \rangle \geq -[(1-t)\lambda_\gamma(K_0) + t\lambda_\gamma(K_1)]v_t,$$

in  $K_t$ .

$$\Delta v_t - \langle \nabla v_t, z \rangle \geq -[(1-t)\lambda_\gamma(K_0) + t\lambda_\gamma(K_1)]v_t$$

$$\Delta v_t - \langle \nabla v_t, z \rangle \geq -[(1-t)\lambda_\gamma(K_0) + t\lambda_\gamma(K_1)]v_t$$

Multiply both sides by  $v_t$ ,

$$\Delta v_t - \langle \nabla v_t, z \rangle \geq -[(1-t)\lambda_\gamma(K_0) + t\lambda_\gamma(K_1)]v_t$$

Multiply both sides by  $v_t$ , integrate on  $K_t$  with respect to  $\gamma_n$ ,



$$\Delta v_t - \langle \nabla v_t, z \rangle \geq -[(1-t)\lambda_\gamma(K_0) + t\lambda_\gamma(K_1)]v_t$$

Multiply both sides by  $v_t$ , integrate on  $K_t$  with respect to  $\gamma_n$ , use the integration by parts formula,

$$\Delta v_t - \langle \nabla v_t, z \rangle \geq -[(1-t)\lambda_\gamma(K_0) + t\lambda_\gamma(K_1)]v_t$$

Multiply both sides by  $v_t$ , integrate on  $K_t$  with respect to  $\gamma_n$ , use the integration by parts formula,

$$\int_{K_t} |\nabla v_t|^2 d\gamma_n \leq [(1-t)\lambda_\gamma(K_0) + t\lambda_\gamma(K_1)] \int_{K_t} v_t^2 d\gamma_n.$$

$$\Delta v_t - \langle \nabla v_t, z \rangle \geq -[(1-t)\lambda_\gamma(K_0) + t\lambda_\gamma(K_1)]v_t$$

Multiply both sides by  $v_t$ , integrate on  $K_t$  with respect to  $\gamma_n$ , use the integration by parts formula,

$$\int_{K_t} |\nabla v_t|^2 d\gamma_n \leq [(1-t)\lambda_\gamma(K_0) + t\lambda_\gamma(K_1)] \int_{K_t} v_t^2 d\gamma_n.$$

Then

$$(1-t)\lambda_\gamma(K_0) + t\lambda_\gamma(K_1) \geq \frac{\int_{K_t} |\nabla v_t|^2 d\gamma_n}{\int_{K_t} v_t^2 d\gamma_n}$$

$$\Delta v_t - \langle \nabla v_t, z \rangle \geq -[(1-t)\lambda_\gamma(K_0) + t\lambda_\gamma(K_1)]v_t$$

Multiply both sides by  $v_t$ , integrate on  $K_t$  with respect to  $\gamma_n$ , use the integration by parts formula,

$$\int_{K_t} |\nabla v_t|^2 d\gamma_n \leq [(1-t)\lambda_\gamma(K_0) + t\lambda_\gamma(K_1)] \int_{K_t} v_t^2 d\gamma_n.$$

Then

$$(1-t)\lambda_\gamma(K_0) + t\lambda_\gamma(K_1) \geq \frac{\int_{K_t} |\nabla v_t|^2 d\gamma_n}{\int_{K_t} v_t^2 d\gamma_n} = R_\gamma[v_t]$$

$$\Delta v_t - \langle \nabla v_t, z \rangle \geq -[(1-t)\lambda_\gamma(K_0) + t\lambda_\gamma(K_1)]v_t$$

Multiply both sides by  $v_t$ , integrate on  $K_t$  with respect to  $\gamma_n$ , use the integration by parts formula,

$$\int_{K_t} |\nabla v_t|^2 d\gamma_n \leq [(1-t)\lambda_\gamma(K_0) + t\lambda_\gamma(K_1)] \int_{K_t} v_t^2 d\gamma_n.$$

Then

$$\begin{aligned} (1-t)\lambda_\gamma(K_0) + t\lambda_\gamma(K_1) &\geq \frac{\int_{K_t} |\nabla v_t|^2 d\gamma_n}{\int_{K_t} v_t^2 d\gamma_n} = R_\gamma[v_t] \\ &\geq \lambda_\gamma((1-t)K_0 + tK_1). \end{aligned}$$

$$\Delta v_t - \langle \nabla v_t, z \rangle \geq -[(1-t)\lambda_\gamma(K_0) + t\lambda_\gamma(K_1)]v_t$$

Multiply both sides by  $v_t$ , integrate on  $K_t$  with respect to  $\gamma_n$ , use the integration by parts formula,

$$\int_{K_t} |\nabla v_t|^2 d\gamma_n \leq [(1-t)\lambda_\gamma(K_0) + t\lambda_\gamma(K_1)] \int_{K_t} v_t^2 d\gamma_n.$$

Then

$$\begin{aligned} (1-t)\lambda_\gamma(K_0) + t\lambda_\gamma(K_1) &\geq \frac{\int_{K_t} |\nabla v_t|^2 d\gamma_n}{\int_{K_t} v_t^2 d\gamma_n} = R_\gamma[v_t] \\ &\geq \lambda_\gamma((1-t)K_0 + tK_1). \end{aligned}$$





Salzburg, 2004